

**SALMON SPAWNING GROUND SURVEYS
IN THE BRISTOL BAY AREA, ALASKA, 1997**



by

**James B. Browning, Jeffrey R. Regnart,
and Keith A. Weiland**

Regional Information Report¹ No. 2A98-34

**Alaska Department of Fish and Game
Division of Commercial Fisheries
333 Raspberry Road
Anchorage, Alaska 99518-1599**

September 1998

¹ The Regional Information Report Series was established in 1987 to provide an information access system for all unpublished divisional reports. These reports frequently serve diverse ad hoc informational purposes or archive basic uninterpreted data. To accommodate timely reporting of recently collected information, reports in this series undergo only limited internal review and may contain preliminary data; this information may be subsequently finalized and published in the formal literature. Consequently, these reports should not be cited without prior approval of the author or the Division of Commercial Fisheries Management and Development.

AUTHORS

James Browning is the Nushagak Area Management Biologist for the Alaska Department of Fish and Game, Commercial Fisheries Division, P.O. Box 230, Dillingham, AK 99576-0230.

Jeffrey R. Regnart is the Naknek/Kvichak Area Management Biologist for the Alaska Department of Fish and Game, Commercial Fisheries Division, 333 Raspberry Rd, Anchorage, AK 99518.

Keith A. Weiland is the Egegik and Ugashik Area Management Biologist for the Alaska Department of Fish and Game, Commercial Fisheries Division, 333 Raspberry Road, Anchorage, AK 99518.

ACKNOWLEDGMENTS

We would like to thank the U.S. Fish and Wildlife Service and the University of Washington, Fisheries Research Institute for equipment, personnel and funding they provided to help gather escapement data in 1997. Critical review of this manuscript by Stephen M. Fried is gratefully acknowledged.

TABLE OF CONTENTS

	<u>Page</u>
LIST OF TABLES	iv
LIST OF FIGURES	vi
LIST OF APPENDICES	vii
INTRODUCTION	1
Naknek/Kvichak District	1
Egegik District	1
Ugashik District	2
Nushagak District	2
Togiak District	4
METHODS	4
Naknek/Kvichak District	5
Egegik District	5
Ugashik District	5
Nushagak District	6
Togiak District	6
RESULTS AND DISCUSSION	7
Naknek-Kvichak District	7
Egegik District	8
Ugashik District	9
Nushagak District	10
Togiak District	10
LITERATURE CITED	12
TABLES	14
FIGURES	30
APPENDICES	40

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1. Aerial survey counts of sockeye salmon, Alagnak River system, 1997	14
2. Aerial survey counts of chinook, chum, pink, and coho salmon, Naknek-Kvichak District, 1997	15
3. Aerial survey peak counts of chinook salmon escapement, Egegik District, 1997	16
4. Aerial survey peak counts of chum salmon escapement, Egegik District, 1997	17
5. Aerial survey peak counts of coho salmon escapement, Egegik District, 1997	18
6. Aerial survey peak counts of sockeye salmon escapement, King Salmon and Dog Salmon Rivers, Ugashik District, 1997	19
7. Peak survey counts of chinook salmon escapement, Ugashik District, 1997	20
8. Peak survey counts of chum salmon escapement, Ugashik District, 1997	21
9. Aerial survey peak counts of coho salmon escapement, Ugashik District, 1997	22
10. Peak aerial counts of live sockeye salmon and total escapement estimates, Wood River system, 1997	23
11. Peak aerial counts of live sockeye salmon and total escapement estimates, Togiak District, 1997	26
12. Peak aerial counts of live chinook salmon and total escapement estimates, Togiak District, 1997	27

LIST OF TABLES (Continued)

<u>Table</u>	<u>Page</u>
13. Peak aerial counts of live chum salmon and total escapement estimates, Togiak District, 1997	28
14. Peak aerial counts of live coho salmon and total escapement estimates, Togiak District, 1997	29

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1. Bristol Bay management area, Alaska	30
2. Alagnak River drainage, Bristol Bay, Alaska	31
3. Egegik River drainage, Bristol Bay, Alaska	32
4. Ugashik River drainage, Bristol Bay, Alaska	33
5. Wood River Lakes system, Bristol Bay, Alaska	34
6. Lake Nunavaugaluk system, Bristol Bay, Alaska	35
7. Nushagak-Mulchatna River system, Bristol Bay, Alaska	36
8. Tikchik Lakes system, Bristol Bay, Alaska	37
9. Togiak River system, Bristol Bay, Alaska	38
10. Kulukak River system, Bristol Bay, Alaska	39

LIST OF APPENDICES

<u>Table</u>	<u>Page</u>
1. Sockeye salmon total escapement estimates, Naknek-Kvichak District, 1955-1997.....	40
2. Aerial survey counts of chinook salmon escapements, Naknek River drainage, 1970-1997.....	42
3. Chinook salmon escapement survey history, mainstem Naknek River, 1929-1997	43
4. Chinook salmon escapement survey history, Big Creek, Naknek River drainage, 1963-1997	46
5. Chinook salmon escapement survey history, King Salmon Creek, Naknek River drainage, 1964-1997	49
6. Chinook salmon escapement survey history, Paul's Creek, Naknek River drainage, 1971-1997	51
7. Chinook salmon escapement survey history, Alagnak River, 1963-1997	53
8. Chinook salmon escapement survey history, Kvichak River, 1932-1997	55
9. Chinook salmon escapement data, Naknek-Kvichak District, 1970-1997	56
10. Chum salmon escapement survey history, Alagnak River, 1961-1997	57
11. Pink salmon escapement survey history, Alagnak River, 1968-1997	59
12. Pink salmon escapement survey history, Kvichak River, 1966-1997	60
13. Pink salmon escapement survey history, Naknek River, 1974-1997	61
14. Aerial survey counts of chinook salmon escapement, Egegik District, 1981-1997	62
15. Aerial survey counts of chum salmon escapement, Egegik District, 1982-1997	63

LIST OF APPENDICES (Continued)

<u>Table</u>	<u>Page</u>
16. Aerial survey counts of pink salmon escapement, Egegik District, 1974-1997	64
17. Aerial survey counts of coho salmon escapement, Egegik District, 1981-1997	65
18. Aerial survey counts of chinook salmon escapement, Ugashik District, 1980-1997	66
19. Aerial survey counts of chum salmon escapement, Ugashik District, 1980-1997	67
20. Aerial survey counts of pink salmon escapement, Ugashik District, 1980-1997	68
21. Aerial survey counts of coho salmon escapement, Ugashik District, 1981-1997	69
22. Spawner distribution and total escapement estimates of sockeye salmon, Wood River system, 1959-1997	70
23. Total escapement estimates of pink salmon, Nushagak and Togiak Districts, 1962-1997	71
24. Aerial estimates of sockeye salmon escapements, Togiak District 1977-1997	72
25. Peak aerial counts of live sockeye salmon, Togiak River drainage, 1977-1997	73
26. Peak aerial counts of live sockeye salmon, Togiak District, 1977-1997	74
27. Peak aerial counts of live chinook salmon, Togiak River drainage, 1977-1997	75
28. Peak aerial counts of live chinook salmon, Togiak District, 1977-1997	76
29. Peak aerial counts of live chum salmon, Togiak River drainage, 1977-1997	77

LIST OF APPENDICES (Continued)

<u>Table</u>	<u>Page</u>
30. Peak aerial counts of live chum salmon, Togiak District, 1977-1997	78
31. Peak aerial counts of live coho salmon, Togiak River drainage, 1980-1997	79
32. Peak aerial counts of live coho salmon, Togiak District, 1980-1997	80

INTRODUCTION

Aerial surveys of salmon spawning streams have been conducted in the Bristol Bay area of Alaska (Figure 1) for many years to provide biologists with information regarding the abundance and distribution of sockeye salmon *Oncorhynchus nerka*, chinook salmon *O. tshawytscha*, chum salmon *O. keta*, pink salmon *O. gorbuscha*, and coho salmon *O. kisutch* escapements. This information is important to fishery managers for several reasons. It supplements data gathered at counting towers on the mainstem rivers, provides data from rivers where counting towers are not utilized, and provides data for time periods and species not covered by counting tower operations. Data collected is used to: (1) evaluate escapement goals and escapement/return relationships, (2) forecast future returns, (3) identify possible management problems relating to escapements, and (4) contribute to strategies designed to alleviate escapement problems. This report summarizes the 1997 salmon spawning ground surveys conducted in the Bristol Bay area.

Naknek/Kvichak District

Naknek-Kvichak District is comprised of three major rivers: (1) the Kvichak River, issuing from Iliamna Lake and its tributaries, (2) the Alagnak or Branch River flowing from Kukaklek and Nonvianuk Lakes, and (3) the Naknek River emanating from Naknek Lake and its tributaries (Figure 2). All of these systems flow into Kvichak Bay.

Since 1955, Kvichak River sockeye salmon escapements have been estimated using counting towers located on the mainstem river, approximately one quarter mile downstream of Lake Iliamna's outlet. From 1957 to 1976, Alagnak River sockeye escapements were estimated using a counting tower located near the upper extent of tidal influence. Since 1977, all Alagnak sockeye escapements have been estimated using aerial surveys. From 1950 to 1957, sockeye escapements to the Naknek River system were counted using a weir on the mainstem river just upstream of the tidal influence. From 1958 to the present, escapements have been estimated using counting towers near the Naknek River 'Rapids' downstream of the outlet of Naknek Lake. Escapement of other salmon species into Naknek-Kvichak District drainages have been estimated using aerial surveys.

Egegik District

Egegik River system contains two major watersheds: (1) the Egegik River, emanating from Becharof Lake and nearby coastal lowlands, and (2) the King Salmon River, issuing from runoff from the Kejulik Mountains and southern portions of Katmai National Park (Figure 3). Both rivers flow into Egegik Bay near Egegik village.

From 1952 through 1956, a weir was used in the Egegik River to count sockeye salmon escapements. The weir was located near the base of the Egegik River 'Rapids'. From 1957 to the present, counting towers situated between the outlet of Becharof Lake and Egegik Lagoon have been used to estimate sockeye escapements. Escapements for other salmon species have been estimated using aerial surveys.

Ugashik District

The Ugashik River system is comprised of four major watersheds: (1) the Ugashik River, flowing from Lower Ugashik Lake and nearby coastal lowlands, (2) the Dog Salmon River, emanating from glacial melt and runoff from peaks in the Aleutian Range, (3) the King Salmon River, issuing from Mother Goose Lake and three major runoff tributaries, and (4) Dago Creek, emitting from a large lowland coastal area (Figure 4). All of these systems flow into the intertidal reaches of Ugashik River and Ugashik Bay.

From 1949 to 1956, a weir located downstream from the outlet of Lower Ugashik Lake was used to count sockeye salmon escapements. From 1957 to the present, sockeye escapements have been estimated using counting towers located between the outlet of Lower Ugashik Lake and Ugashik Lagoon. Escapements for other salmon species have been estimated using aerial surveys.

Nushagak District

Nushagak watershed is comprised of four major rivers: (1) the Wood River, draining Grant, Kulik, Beverley, Nerka, and Aleknagik Lakes, (2) the Nushagak River, draining Tikchik Lakes and the Nuyakuk, upper Nushagak, and Mulchatna Rivers, (3) the Igushik River, draining Ualik and Amanka Lakes, and (4) the Snake River, draining Lake Nunavaugaluk (Figures 5 through 8). All of these systems empty into Nushagak Bay.

Abundance and age composition of sockeye salmon escapements in the Wood River Lake system has been estimated annually from counting towers at the outlet of Lake Aleknagik since 1953.

Sockeye salmon distribution in the Wood River Lake system is an important element in establishing escapement goals and measuring success in achieving escapement goals for this system. Interconnecting rivers between the large lakes in the system are primarily used by three-ocean sockeye for spawning, while the lake beaches and tributary streams are used more by two-ocean sockeye. Knowledge of the age composition of returning sockeye gives managers the ability to use a variable escapement goal policy to minimize overcrowding of spawners in the interconnecting rivers while taking advantage of the extensive beach spawning areas and numerous tributary streams.

ADF&G staff have conducted aerial surveys to assess sockeye spawner distribution within the Wood River Lake system each year. Personnel from the University of Washington, Fisheries Research Institute also conduct ground surveys on major creeks and some rivers of the system. Surveys of the actual spawning distribution within the creeks, rivers, and beaches of the system provide a measure of management success in obtaining the desired spawning distribution.

Salmon escapement in the Nushagak River is estimated by a sonar project, located on the Nushagak River below Portage Creek, approximately 32 km (20 miles) upstream from the river mouth. The Nushagak River sonar project has been used since 1980 to estimate annual escapements for all salmon species in the entire Nushagak drainage (Miller 1997). Prior to the advent of the sonar

project, sockeye escapement was estimated by a counting tower project on the Nuyakuk River (1959-1988). Aerial surveys of the Nushagak-Mulchatna system were conducted annually beginning in 1966. Initial surveys provided escapement estimates for chinook and chum salmon, and surveys in the Nushagak and Mulchatna systems since 1977 were used to estimate sockeye abundance in that system. Together, the combined estimates from counting towers and aerial surveys were used by fishery managers as estimates of the Nushagak River drainage sockeye escapement.

ADF&G staff continued to survey the upper Nushagak and Mulchatna areas after the development of the sonar project to provide a comparison with sonar estimates and document spawner distribution for all species except coho salmon. Chum salmon surveys were discontinued in the Nushagak District in 1980, and surveys of the Nushagak-Mulchatna Rivers for all other species were discontinued in 1991 due to the success of the sonar project and limited funding. After terminating the Nuyakuk tower project in 1988, and terminating surveys of the Nushagak and Mulchatna systems in 1991, little information was available to assess sockeye spawning distribution in the Nushagak River.

Aerial surveys were conducted sporadically in the Tikchik Lakes system from 1954 to 1987 to assess spawner distribution of sockeye salmon. Surveys of the Tikchik Lakes were conducted sporadically since 1990 to document an apparent change in spawner distribution, evidenced by changes observed in the age composition of Nushagak River sockeye escapement, and supported by reports of low numbers of spawners in the Tikchik Lake system. These surveys have documented lower than expected numbers of spawners in the Tikchik Lakes system, based on sonar estimates in the lower Nushagak River and historical distribution patterns (Brookover et. al. 1995). However, few corresponding surveys were conducted in the Nushagak and Mulchatna drainages to completely assess distribution. With the assistance of Fisheries Research Institute personnel, the counting tower project on the Nuyakuk River was re-initiated beginning in 1995 in order to assess recent distribution and production trends in the Nushagak drainage.

Sockeye escapement is measured in the Igushik Lakes system at a counting tower located at the outlet of Amanka Lake. Spawner distribution has not been documented annually, and surveys have not been conducted on the Igushik system for sockeye salmon and other species since 1991 (Russell, et. al. 1992). Spawning escapement and distribution of sockeye salmon in the Snake Lake system was estimated annually by aerial surveys, but funding was not available for these surveys from 1991 through 1994.

Togiak District

Togiak District includes two major river drainages: (1) the Togiak River, draining Togiak, Gechiak, Pungokebuk, and Ongivinuck Lakes and Naylorun and Kemuk Rivers (Figure 9), and (2) the Kulukak River, draining Kulukak Lake (Figure 10). Various smaller systems within the district include the Tithe Creek Ponds and the Quigmy, Matogak, Osviak, Slug, Negukthlik, and Ungalikthluk Rivers. Kulukak River and the Tithe Creek Ponds flow into Kulukak Bay, located in the eastern portion of the district; the Togiak and Quigmy Rivers flow into Togiak Bay, located in the middle of the district; and the Matogak, Osviak, and Slug Rivers flow into Hagemeister Straits and coastal waters in the western portion of the district (Figure 1).

Sockeye salmon escapement is estimated for the Togiak Lake system from counting towers operated at the outlet of Togiak Lake. Abundance and distribution of spawning populations of sockeye salmon in the Togiak River and tributaries below the counting towers, as well as other systems within the Togiak District, are estimated by aerial surveys. Abundance and distribution of chinook, chum, pink, and coho salmon spawning in Togiak District watersheds are also estimated entirely from aerial surveys.

Since 1991, the operational budget has not had sufficient funds to conduct spawning ground aerial surveys in the Togiak District. However, the USFWS and Togiak National Wildlife Refuge has provided some additional funding for aircraft charters for aerial surveys in the District to monitor salmon populations on the refuge drainages.

METHODS

All survey flights were conducted from small fixed-wing, high-wing, wheeled aircraft (Super Cub, Cessna 180, Cessna 185, or Cessna 206) or helicopter (Robinson R-22) chartered from local air charter companies and flown by experienced survey pilots. Several surveys in the Togiak National Wildlife Refuge were flown by USFWS pilots and aircraft. Salmon were counted by Alaska Department of Fish and Game (ADF&G) or USFWS biologists familiar with the streams and target species. Counts were made from low altitudes (200 to 400 feet) at air speeds of 50 to 90 mph. Polarized sunglasses and aircraft positioning were used to minimize effects of glare off the water. Surveys were scheduled to coincide as closely as possible to the historic peak of spawning for the target species, taking into account weather, water conditions, and aircraft availability. Peak of spawning was defined as that point when the greatest number of spawning salmon are occupying redds. Counts were registered on a hand tally counter or on a tape player. This information was transferred to survey data forms either sometime during the survey or upon returning to the office.

Aerial surveys account for only a portion of the known spawning populations (Evzerof, 1975; Nielson and Green, 1981; Rogers, 1984). At the time of each survey, some of the salmon have yet to reach the spawning grounds, some have already spawned and died, some are still schooled, and some are either misidentified or not seen. Methods used to interpret aerial survey counts are described below for each commercial fishing district.

Naknek/Kvichak District

Aerial surveys were flown during late summer and fall to assess escapements of sockeye, chinook, and chum salmon in portions of the Naknek-Kvichak District. Salmon counts for these drainages are indices of the total number of each species present in the spawning area at the time of the survey. Two surveys were flown, August 7 and 16, to provide estimates of Alagnak River drainage sockeye, chinook, and chum escapements. Additionally, all major chinook spawning areas in the Naknek River drainage were surveyed between July 18 and August 16, and the Kvichak River chinook escapement survey was flown August 15. These survey counts were not expanded to provide instantaneous population estimates, although expansions have been made in some earlier years based on subjective criteria.

Counting towers were used to estimate total sockeye salmon escapements to the Kvichak and Naknek Rivers. A late summer survey of sockeye salmon spawning distribution in the Kvichak River system was completed August 15, 16, and September 4, and the results were documented in Regnart (1996). All aerial survey counts in the district were made by ADF&G, Commercial Fisheries Management and Development Division staff.

Egegik District

No system-wide aerial surveys were flown for sockeye salmon in 1997. Aerial surveys of all known chinook and chum salmon spawning areas in both the Egegik and King Salmon Rivers were flown on July 27. With funding provided by the Alaska Department of Fish and Game and the U.S. Fish and Wildlife Service (USFWS), aerial surveys were flown on August 18, September 12 and 30, and on October 17 to estimate coho salmon escapements. In addition, a weir was operated on Gertrude Creek by USFWS from April 18 to September 17. All aerial survey counts in the Egegik District reflect only the actual numbers of salmon sighted and should be considered an index of abundance only.

Ugashik District

Salmon counts in the Ugashik District reflect the actual numbers of salmon sighted on the spawning grounds for 1997. Aerial surveys of all known chinook and chum salmon spawning areas in the Ugashik drainage were flown on August 6. With funding provided by the Alaska Department of Fish and Game, aerial surveys were flown on September 30 and October 17 to estimate coho salmon escapements. Aerial survey counts should be considered only an index of total abundance.

Nushagak District

Aerial surveys were conducted to assess spawning distribution of sockeye salmon in the Wood River system in 1997. Survey methods and data analysis for the Nushagak District were similar to those described by Nelson (1979), Bucher (1981), and Russell, Bill and Bucher (1990).

Sockeye salmon escapements for each spawning stream, beach, or river in the Wood River have been estimated using the proportion of sockeye salmon observed at a given location in relation to the tower count. Different expansion factors were assigned to each type of spawning habitat. For a more detailed description of the analysis of Wood River survey counts, see Nelson (1973).

Togiak District

Survey and data analysis methods used in the Togiak District were similar to those described by Nelson (1979), Bucher (1981), and Russell, et. al. (1990). Aerial surveys of spawning sockeye, chinook, chum, and coho salmon were conducted at the peak of spawning for each species, using criteria similar to Nelson (1979) and Bucher (1981). Survey coverage was provided primarily by ADF&G staff, with some counts provided by USFWS Togiak National Wildlife Refuge staff.

Peak aerial survey counts for sockeye salmon in the Togiak Lake system above the counting tower have generally accounted for 47% (range: 40% - 50%) of the escapements estimated at the tower (Nelson 1967). Therefore, peak aerial counts of sockeye salmon in systems without counting towers (i.e. Kulukak River, mainstem and tributaries of the Togiak River below the towers) were multiplied by 2.0 to estimate total escapement. Since 1980, aerial counts of chinook salmon in the Togiak District have generally been multiplied by 2.5 to estimate total escapement. During the 1997 surveys, a 1.5 factor was applied to the mainstem and tributaries of the Togiak River due to low and extremely clear water, while a factor of 2.0 was used in some of the smaller systems in the western portion of the district. Since 1968, aerial counts of chum salmon have generally been multiplied by 2.0 (Nelson 1968). Since 1978, pink salmon escapements have also been estimated by multiplying aerial counts by 2.0. An expansion factor of 3.0 has been used for coho salmon in all areas of the Togiak District since the initiation of coho surveys in 1980. Expansion factors have been subjectively adjusted based on weather conditions, visibility, and survey timing with respect to the peak spawning activity.

Aerial counts of sockeye salmon in the Slug, Matogak, Osviak, Negukthlik, and Ungalikthluk rivers were obtained incidentally during chum and chinook salmon escapement surveys, following the methods used by Nelson (1979).

RESULTS AND DISCUSSION

Naknek/Kvichak District

Aerial surveys of sockeye salmon escapement into the Alagnak River and its tributaries were conducted on August 7 and 16. The sockeye salmon escapement index count totaled 218,115 for this system (Table 1). This count was slightly above the average (1977-1994) aerial count of 204,000, and was approximately 18% greater than the escapement point goal of 185,000.

Aerial surveys of chinook salmon escapements into the Naknek River drainage were flown from July 18 through August 16. Chinook salmon escapement counts were made in each of the four main spawning areas: mainstem Naknek River, Big Creek, King Salmon Creek, and Paul's Creek. A total of 10,453 chinook salmon were counted. The largest components of this total were counts of 1,783 chinook in Big Creek on August 7, and 7,520 chinook in the mainstem Naknek River on August 16 (Table 2). Over the period from 1970-1996 there have been 17 years in which chinook salmon escapement indices have been obtained from all four main spawning areas (Appendix Tables 2-6). The chinook escapement index for these 16 years has ranged from a low of 2,691 in 1992 to a high of 11,730 in 1988.

Alagnak River drainage chinook salmon escapement was surveyed on August 7, yielding a count of 15,210 (Table 2). From 1970-1996, Alagnak chinook salmon counts have ranged from a low of 824 in 1973 to a high of 11,650 in 1978. An aerial survey of chinook salmon escapement into the Kvichak River was conducted on August 15 and yielded a count of 103.

Chum salmon were counted only during the August 7 Alagnak River aerial surveys. Alagnak River has been the principal chum salmon producing system in the Naknek-Kvichak District. A total of 37,800 spawning chum salmon were observed during the 1997 survey.

Egegik District

The 1997 Egegik River sockeye escapement past the counting towers totaled 1,103,964 fish, or 10% above the point goal (1.0 million) but below the upper limit of the BEG range of 800 thousand to 1.4 million. Forty additional sockeye salmon were observed on a post season aerial survey.

Aerial survey peak counts of all known chinook spawning areas in the Egegik drainage yielded a total count of 1,616 chinook salmon (Table 3). An additional 30 chinook were counted past Egegik River counting towers bringing the district escapement index to 1,646 fish. This total was the second highest on record and is only superseded by a helicopter survey in 1994 (Appendix Table 14). The commercial chinook harvest in the Egegik District totaled approximately 2,143 fish, 27% below the 1977-1996 average harvest of 2,926. Prohibiting usage of gillnets with greater than 5.5 inch mesh, from June 1-July 1 facilitated passage of chinook salmon through the commercial fishing district, but good catches between June 10 and June 23 suggested the chinook run was perhaps stronger than normal. Given the catch and escapement figures above, the Egegik chinook salmon removal rate for 1997 was around 57%.

The peak chum salmon escapement index was obtained July 27 and totaled 1,723 fish (Table 4). The 1997 index was well below the 1982-1996 average of 9,474 fish (Appendix Table 15). The 1997 commercial chum harvest from the Egegik District totaled approximately 53,000 fish, 49% below the 1977 to 1996 average of 103,000. Though escapement indices of less than 10,000 chum salmon have been recorded in the district in each of the last nine years, it is believed that aerial surveys for chum salmon are not a reliable indicator of chum escapements. For example, the weir operated by USFWS on Gertrude Creek counted over 11,000 chum salmon while the peak aerial count for that system was around 500 fish, or 22 times less than the weir count. Given this amount of discrepancy in only one tributary, it is likely that chum escapement indices documented over the last several years, by airplanes, have probably greatly under estimated true chum escapements.

No pink salmon were noted during the July 27 aerial survey. Two pink salmon were reported from the commercial catch. The 1974 to 1997 pink salmon escapement indices are presented in Appendix Table 16.

The coho salmon escapement was documented with aerial surveys conducted on August 18, September 12 and 30, and October 17 (Table 5). Funding for these surveys was provided by the U.S. Fish and Wildlife Service and the Alaska Department of Fish and Game. A combined total of 4,106 coho salmon were counted in the King Salmon and Egegik Rivers and in numerous tributaries of Becharof Lake. Of this total, approximately 3,455 fish were counted upstream of the Egegik River counting towers. The aerial counts were focused on the main coho salmon producing areas. Compared to the last two years, counts in most of the areas surveyed were below the 2-year averages. The commercial harvest totaled approximately 36,000 fish, which was about equal to the 1977 to 1996 average of 36,400. Deliveries occurred through September 4, though officially the fishery was open until September 30. Historical survey counts are listed in Appendix Table 17.

Ugashik District

The 1997 sockeye salmon escapement past Ugashik River counting tower was approximately 618,400 fish, or 12% below the point goal of 700,000. No system-wide aerial surveys were conducted because of a lack of funding, however; an additional 10,600 and 27,600 sockeye salmon were counted in the Dog Salmon and King Salmon Rivers, respectively, during a chinook and chum salmon survey (Table 6).

Chinook salmon escapement surveys of Dog Salmon, King Salmon, and Ugashik Rivers were flown on August 6 and yielded a count of 4,022 fish. Additionally, 144 chinook salmon were counted past the counting towers bringing the cumulative escapement count to 4,166 (Table 7). The Dog Salmon River count of 906 was the largest escapement component for the system. The 1997 escapement count was 11% below the 1980 to 1996 average count of 4,696 chinook salmon (Appendix Table 18). The Ugashik District's commercial catch of 1,096 chinook was 72% below the average harvest of 3,900. Overall, the 1997 Ugashik chinook run was above average.

Aerial surveys of Dog Salmon, King Salmon, and Ugashik Rivers on August 6, yielding a count of 30,115 chum salmon. An additional 30 chum salmon were counted past the counting towers bringing the escapement total to 30,145 (Table 8). The survey was considered to be prior to the peak of spawning as most fish were still schooled up. The 1997 aerial count was 7% below the 1980-1996 average of 32,600 (Appendix Table 19). The District's commercial chum salmon harvest was approximately 16,400 fish. The catch was well below the 1977 to 1996 average of 71,200.

The Ugashik pink salmon returns have historically been very small, and this year, again, was no exception. No pink salmon were reported in the commercial catch. However, fishing only a total of 55 hours during the whole EO period, from June 23 to July 27 was most likely a reason for the lack of pink salmon, as well as chinook and chum salmon, in this year's harvest. No pink salmon were observed on the escapement survey flown on August 6, nor were any counted past the tower before it ceased operation on July 29 (Appendix Table 20).

Aerial surveys for coho salmon were again made this year in the Ugashik drainage and are listed in Table 9. A total of 9,400 were observed most of which came from the King Salmon River. Daily commercial coho catch statistics for set net gear ranged below to above average and the district remained on normal fishing time except for a reduction during one week of the season. The coho harvest of approximately 7,600 fish was well below average. Historic coho salmon escapement data is recorded in Appendix Table 21.

Nushagak District

Spawning ground surveys for chinook salmon have been flown sporadically in the Nushagak District for the last several years due to budgetary shortfalls. The Portage Creek sonar counting project produced an apportioned estimate of 40,705 chinook salmon in the Nushagak River for 1997. Due to low flows and elevated water temperatures in 1997, fish migration routes and position in the water column relative to the sonified area was suspect. An aerial survey flown on July 21 documented approximately 42,000 chinook salmon in the index areas historically surveyed. This figure expanded for total spawning area became an estimate of 55,000 chinook and further expanded for visibility factor yielded an estimate of 82,000 spawning chinook salmon for 1997.

Spawning escapement of sockeye salmon in the Wood River system was estimated to be 1.51 million fish, 51% greater than the escapement goal. Two-ocean sockeye comprised approximately 56% of the Wood River escapement while three-ocean sockeye contributed another 42% of the escapement, hence Wood River was managed for the 1.2 million end of the escapement goal range. Poor sockeye escapement in the Nushagak River prompted conservative management of the Nushagak District commercial fishery in 1997, and conservation concerns for coho salmon prompted a closure in Nushagak District for coho salmon on July 23 as directed by the Nushagak River Coho Salmon Management Plan. The combination of these management actions was the primarily reason for the large escapement in the Wood River system in 1997. The Wood River Special Harvest Area remained open through July 29 to harvest additional sockeye salmon surplus to the escapement goal.

Aerial surveys were flown on July 28, August 4, 16 and 28 on the Wood River Lakes drainage. Spawning sockeye salmon were observed to be predominantly distributed in riverine habitat in 1997 (Table 10, Appendix Table 22). Most spawning was observed on interconnecting rivers, followed by lake beaches and creeks. Also, due to extremely low water levels and flows, access to creek spawning habitat was obstructed by beaver dams to a larger degree than in previous years.

Togiak District

Peak aerial counts and total population estimates were derived from aerial surveys for sockeye salmon in major river systems of the Togiak District in 1997. (Table 12) The expanded aerial survey estimate of 20,625 sockeye salmon for the Togiak River and its tributaries below the counting tower was 22% below the 1977-1996 average of 26,632 fish (Appendix Table 24). Escapement past the counting tower was 131,682 sockeye, 12% under the escapement goal of 150,000. The spawning escapement of sockeye salmon in Kulukak Section, including Kulukak River, Kulukak Lake, and Tithe Creek Ponds, was estimated to be 7,950 fish, only 28% of the 1987-1996 average of 27,534. Peak aerial sockeye salmon counts into the mainstem portion of the Togiak River were above the 1977-1996 average (Appendix Tables 25 and 26), while counts for all of the tributaries to the Togiak River were below average. Total sockeye salmon escapement for Togiak District was 166,627 fish, with the Quigmy and Ungalikthluk Rivers uncounted. Due to weather, surveys were conducted after the peak of spawning on several

systems, while the majority of systems were surveyed at peak spawning levels in the Togiak and Kulukak River drainages.

Aerial surveys for peak live counts and expanded escapement estimates for chinook salmon were conducted in all the major drainages within the Togiak District for 1997 (Table 12). The minimum expanded escapement estimate for Togiak District was 11,495 chinook salmon, with the Ungalikthluk and Negukthlik Rivers remaining uncounted. The 1997 district-wide chinook escapement was 22% below the 1992-1996 average of 14,743 fish (ADF&G, 1997). The aerial peak live count for the Togiak River and tributaries was 19% above the 1977-1996 average, and aerial counts for all chinook systems in the Togiak District combined were 10% below the 20-year average (Appendix Tables 27 and 28). The escapement goal of 10,000 chinook salmon into the Togiak River was achieved with a final expanded estimate of spawning escapement of 10,300 fish. Chinook peak aerial counts for smaller river systems within the district were generally all below average. Kulukak River escapement estimate (775 fish) was poor and comprised only 29% of the 1977-1996 average.

Conditions and timing were fair to poor for most of the chinook salmon systems surveyed. A standard multiplier of 2.5 was applied to most of the aerial counts, however, due to low and clear water conditions in the mainstem of the Togiak River and tributaries, a 2.0 multiplier was used. Surveys for streams west of Togiak River, Negukthlik and Ungalikthluk Rivers were flown in conjunction with USFWS. Spawning activity for chinook appeared to peak on or near August 1 in the Togiak District.

Chum salmon escapement was poor in the entire Togiak District, and was estimated to be 106,580 (Table 13). The 1997 estimate is 55% below the 1977-1996 average (239,489 chum) reported by ADF&G (1997). Peak counts of chum salmon were well below the 1977-1996 average in all streams surveyed within the Togiak District (Appendix Tables 29 and 30).

Chum salmon counts were conducted coincidentally with the chinook salmon surveys. Survey timing was generally post-peak for spawning activity. Significant numbers of chum salmon carcasses were observed in all rivers surveyed. However, a multiplier of 2.0 was still used.

Aerial surveys yielding peak live counts and expanded escapement estimates for coho salmon were successfully completed for most major systems in Togiak District for 1997. Total coho escapement for Togiak River and tributaries was estimated to be 20,625 fish (Table 14). This was 53% below the 1980-1996 average of 43,957 fish. District-wide coho escapement counts were also 53% below the average for the same period (Appendix Tables 31 and 32). Coho salmon appeared to be mostly on spawning beds with few schools observed, and no carcasses were visible, indicating that the survey was probably at or near the peak of spawning activity.

LITERATURE CITED

- ADF&G (Alaska Department of Fish and Game). *In press*. Annual Management Report, 1997, Bristol Bay Area. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division. Anchorage.
- Bucher, W.A. 1981. Spawning ground surveys in the Nushagak and Togiak Districts of Bristol Bay, 1980. Alaska Department of Fish and Game, Division of Commercial Fisheries, Bristol Bay Data Report 81, Anchorage.
- Evzerov, A.V. 1975. An evaluation of the errors occurring in salmon census by aerial survey. In Canadian Translation Fisheries and Aquatic Sciences. No. 4714, 1981. [Transl. from Russian] (Otsenka pogreshnogtei aerovizual'nogo metoda ucheta Lososei), p. 82-84. From Lososevye Dal'Nego Vostoka (CVI:1975).
- Miller, J.X. 1997. Sonar enumeration of pacific salmon escapement into the Nushagak River, 1997. Alaska Department of Fish and Game, Division of Commercial Fisheries Management and Development, Regional Information Report No. 2A96-08, Anchorage.
- Nielson, J.D. and G.H. Green, 1981. Enumeration of spawning salmon from spawner residence time and aerial counts. Transactions of the American Fisheries Society 110:554-556.
- Nelson, M.L. 1967. Red salmon spawning ground surveys in the Nushagak and Togiak Districts, Bristol Bay, 1966. Alaska Department of Fish and Game, Division of Commercial Fisheries, Informational Leaflet 96.
- Nelson, M.L. 1968. Spawning ground surveys in the Nushagak and Togiak Districts, Bristol Bay, 1968. Alaska Department of Fish and Game, Division of Commercial Fisheries, Bristol Bay Data Report 5.
- Nelson, M.L. 1973. Spawning ground surveys in the Nushagak and Togiak Districts of Bristol Bay, 1973. Alaska Department of Fish and Game, Division of Commercial Fisheries, Bristol Bay Data Report 46.
- Nelson, M.L. 1979. Spawning ground surveys in the Nushagak and Togiak Districts of Bristol Bay, 1977-1979. Alaska Department of Fish and Game, Division of Commercial Fisheries, Bristol Bay Data Report 73, Anchorage.

LITERATURE CITED (Continued)

- Regnart, J.R. 1996. Kvichak River sockeye salmon spawning ground surveys, 1955-1996. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report No. 2A95-48, Anchorage.
- Rogers, D.E. 1984. Aerial survey estimates of Bristol Bay sockeye salmon escapements. Proceedings of the Workshop on Stream Indexing for Salmon Escapement Estimation. Canadian Technical Report of Fisheries and Aquatic Sciences. 1326:197-208.
- Russell, R.B., D.L. Bill, and W.A. Bucher. 1990. Salmon spawning ground surveys in the Bristol Bay area, 1989. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 2K89-15, Anchorage.
- Russell, R.B., J.R. Regnart, and T.E. Brookover. 1992. Salmon spawning ground surveys in the Bristol Bay area, 1991. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 2A92-01, Anchorage.
- Regnart, J.R., T.E. Brookover, J.B. Browning and K.A. Weiland. 1996. Salmon spawning ground surveys in the Bristol Bay area, 1996. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 2A97-21, Anchorage.

Table 1. Aerial survey counts of sockeye salmon, Alagnak River system, 1997¹

Location	Number of Fish	Percent of Total
Nonvianuk River	0	0
Nonvianuk Lake	1,350	0.6
Kulik River	45,150	20.7
Kulik Lake	2,650	1.2
Alagnak River	0	0.0
Kukaklek Lake	1,750	0.8
Nanuktuk Creek	27,480	12.6
Battle River	12,650	5.8
Battle Lake	5,300	2.4
Spectacle Creek	85,445	39.2
Funnel Creek	36,340	16.7
Total	218,115	100.0

¹ Aerial surveys were conducted with fixed-wing aircraft.

Table 2. Aerial survey counts of chinook, chum, pink, and coho salmon, Naknek-Kvichak District, 1997.

Location	Survey Date	Number of Salmon			
		Chinook	Chum	Pink	Coho
Kvichak River	15-Aug	103			
Alagnak River	17-Aug	15,210	37,800		
Naknek River :					
Paul's Creek	18-Jul	248	0 ^a		
King Salmon Creek	18-Jul	902	0 ^a		
Big Creek	7-Aug	1,783	0 ^a		
Mainstem Naknek River	16-Aug	10,453			
Total		28,699	37,800		

¹ Aerial surveys were conducted with fixed-wing aircraft.

^a Incidental observation.

Table 3. Aerial survey peak counts of chinook salmon escapement, Egegik District, 1997.

Location	Survey Date	Number of Chinook Salmon Counted
Egegik River	July 17 ^a	30
Shosky Creek	July 27	39
Whale Mountain Creek	July 27 ^b	2
Mossy Creek	July 27	18
Mink Creek	July 27	10
Gertrude Creek	July 27	260
Kaye's Creek	July 27	173
Takayoto Creek	July 27	374
Angle Creek	July 27 ^c	
Contact Creek	July 27	740
Mainstem King Salmon River	July 27 ^c	
Total		1,646

^a Tower count.

^b Creek was mostly dried up with only the lower mile or two watered, but levels were not conducive to fish passage.

^c No counts due to turbid water conditions.

Table 4. Aerial survey peak counts of chum salmon escapement, Egegik District, 1997.

Location	Survey Date	Number of Chum Salmon Counted
Egegik River	July 17 ^a	0
Shosky Creek	July 27	0
Whale Mountain Creek	July 27	220
Mossy Creek	July 27	8
Mink Creek	July 27	10
Gertrude Creek	July 27	495
Kaye's Creek	July 27	290
Takayoto Creek	July 27	60
Angle Creek	July 27 ^b	
Contact Creek	July 27	640
Mainstem King Salmon River	July 27 ^b	
Total		1,723

^a Tower count.

^b No counts due to turbid water conditions.

Table 5. Aerial survey counts of coho salmon escapement, Egegik District, 1997. ^a

Location	Survey Date	Number of Coho Salmon Counted	Comments
<u>Egegik River Drainage¹</u>			
Egegik River Rapids	October 17	1,400	Many schooled off mouth of Myers Creek
Stream 115.8 (Featherly Creek)	September 30	10	
Stream 107.6 (Burl's Creek)	September 30	240	
Stream 90.3 (Salmon Creek)	October 17	750	
Stream 89.8	September 30	10	
Stream 87.0 (Bear Creek)	September 30	60	Includes Margaret & Albert Creeks
Stream 73.5 (Becharof Creek)	September 30	380	
Stream 48.1 (Kejulik River)	September 30	605	
Sub-total		3,455	
<u>King Salmon River Drainage</u>			
Gertrude Creek	September 17	651	U.S. Fish and Wildlife Weir Count
Sub-total		651	
District Total		4,106	

^a Four surveys flown on different dates and only the highest counts are listed.

¹ Streams tributary to Becharof Lake are designated by the number of miles between their mouth and the outlet of Becharof Lake (Egegik River) as one travels around the lake in a clockwise fashion from the Becharof lake outlet. This is the same system of designation used for years by previous investigators. Three surveys were flown on different dates and only the highest counts are listed.

Table 6. Aerial survey peak counts of sockeye salmon escapement, King Salmon and Dog Salmon River, Ugashik District, 1997.

Location	Survey Date	Number of Sockeye Salmon
<u>Ugashik River:</u>		
Grassy Creek	Aug. 6	0
Sub-total		0
<u>King Salmon River System:</u>		
Needle Lake	Aug. 6	600
Volcano Creel	Aug. 6	20
Painter Creek	Aug. 6	27,000
Indecision Creek	Aug. 6	25
Sub-total		27,645
<u>Dog Salmon River System:</u>		
Figure-Eight Creek	Aug. 6	9,100
Goblet Creek	Aug. 6	0
Oldham Creek	Aug. 6	1,300
Wandering Creek	Aug. 6	200
Mainstem Dog Salmon River	Aug. 6	^a
Sub-total		10,600
Total		38,245

^a No fish were observed due to turbid water conditions.

Table 7. Peak survey counts of chinook salmon escapement, Ugashik District, 1997.

Location	Survey Date	Number of Chinook Salmon
<u>King Salmon River System</u>		
Old Creek	Aug. 6	558
Pumice Creek	Aug. 6	536
Painter Creek	Aug. 6	525
Mainstem King Salmon River	Aug. 6	802
Mother Goose Lake	Aug. 6	0
Indecision Creek	Aug. 6	0
Volcano Creek	Aug. 6	0
Sub-total		<u>2,421</u>
<u>Dog Salmon River System</u>		
Figure-Eight Creek	Aug. 6	858
Goblet Creek	Aug. 6	31
Oldham Creek	Aug. 6	17
Wandering Creek	Aug. 6	0
Mainstem Dog Salmon River	Aug. 6	^a
Sub-total		<u>906</u>
<u>Ugashik River System</u>		
Mainstem Ugashik River	Aug. 6	724 ^b
Grassy Creek	Aug. 6	<u>115</u>
Sub-total		839
Total		<u>4,166</u>

^a No fish were observed due to turbid water conditions.

^b Includes tower and below tower counts.

Table 8. Peak survey counts of chum salmon escapement, Ugashik District, 1997.

Location	Survey Date	Number of Chum Salmon
<u>King Salmon River System</u>		
Old Creek	Aug. 6	9,480 ^a
Pumice Creek	Aug. 6	5,300
Painter Creek	Aug. 6	4,200
Mainstem King Salmon River	Aug. 6	10,500
Needle Lake	Aug. 6	115
Indecision Creek	Aug. 6	0
Volcano Creek	Aug. 6	0
Sub-total		29,595
<u>Dog Salmon River System</u>		
Figure-Eight Creek	Aug. 6	430
Goblet Creek	Aug. 6	20
Oldham Creek	Aug. 6	0
Wandering Creek	Aug. 6	0
Mainstem Dog Salmon River	Aug. 6	0 ^b
Sub-total		450
<u>Ugashik River System</u>		
Mainstem Ugashik River	Aug. 6	30
Grassy Creek	Aug. 6	70
Sub-total		100
Total		30,145

^a Includes carcasses.

^b No fish were observed due to turbid water conditions.

Table 9. Aerial survey counts of coho salmon escapement, Ugashik District, 1997. ^a

Location	Survey Date	Number of Coho Salmon Counted	Comments
<hr/>			
<u>Ugashik Drainage</u>			
<u>Upper Ugashik Lake</u>			
Crooked Creek	October 17	970	Upstream
Deer Creek	September 30	380	
 <u>Lower Ugashik Lake</u>			
Black Creek to Cabin	October 17	350	
Black Creek to Elizabeth Lake	October 17	300	
Ugashik Outlet	October 17	250	
 <u>King Salmon River Tributaries</u>			
Pumice Creek	October 17	210	
Old Creek	September 30	580	
Painter Creek	October 17	5,360	
 <u>Dog Salmon River Tributaries</u>			
Figure Eight Creek	October 17	1,000	
<hr/>			
District Total		9,400	

^a Two surveys were flown on different dates and only the highest counts are listed.

Table 10. Peak aerial live counts and total escapement estimates of sockeye salmon in the Wood River system, 1997.

Area	Date	Aerial Count ¹	Population Estimate	Distribution %
Wood River			0	0.0%
Lake Aleknagik		13,025	81,300	5.4%
Eagle Creek				
Hansen Creek				
Happy Creek				
Bear Creek	7/28/97	1,250		
Yako Creek	7/28/97	2,000		
Whitefish Creek	7/28/97	125		
Ice Creek	7/28/97	600		
Mission Creek				
Sunshine Creek	7/28/97	750		
Youth Creek	7/28/97	200		
Northshore Beaches	8/28/97	2,600		
Southshore Beaches	8/28/97	3,800		
Yako Beach	8/28/97	1,700		
Agulowok River & lower River Bay	8/28/97	118,000	176,100	11.6%
Lake Nerka		101,800	635,100	42.0%
Fenno Creek	7/28/97	3,520		
Pike Creek	8/4/97	2,000		
Stovall Creek ²	8/4/97	80		
Bear Creek	8/4/97	0		
Teal Creek	8/4/97	300		
Pick Creek				
Elva Creek				
Kema Creek	8/4/97	3,200		
Hidden Lake Creek	8/4/97	1,000		
Lynx Creek	8/16/97	3,200		
Upper River Bay Beaches, NW	8/28/97	1,600		
Upper River Bay Beaches, SE	8/28/97	5,800		
Allan Cr. - Ross Cr. Beaches	8/28/97	5,500		
N6 - River Bay Beach	8/28/97	800		
Pick Creek Beach	8/28/97	22,100		
Elva Creek Beach	8/28/97	3,700		
Amakuk Arm Beaches	8/28/97	3,600		

(Continued)

Table 10. (Continued)

Area	Date	Aerial Count ¹	Population Estimate	Distribution %
Amakuk Arm - Ott's Bay Beach	8/28/97	4,600		
Ott's Bay Beach	8/28/97	4,100		
Anvil Bay Beaches	8/28/97	13,900		
Anvil Bay - Elbow Pt. Beach	8/28/97	8,300		
Elbow Pt. - Lynx Cr. Beach	8/28/97	6,300		
Lynx Cr. - Teal Cr. Beach	8/28/97	1,200		
Kema Lake Beaches ²				
Hidden Lake Beaches	8/28/97	1,300		
Lynx Lake Beaches	8/28/97	5,700		
Little Togiak River	8/16/97	300	400	0.0%
Little Togiak Lake		9,200	57,400	3.8%
Northshore Beaches	8/28/97	4,300		
Southshore Beaches	8/28/97	4,900		
D Slough Beaches				
Agulukpak River	8/28/97	93,000	138,800	9.2%
Lake Beverley		27,000	168,400	11.1%
Tsun Creek				
Moose Creek	8/14/97	9,300		
Hope Creek	8/14/97	4,600		
Hardluck Bay Beaches	8/28/97	6,600 ^d		
Sam's Beach	8/28/97	600		
Golden Horn Beaches	8/28/97	1,400		
Silver Horn Beaches	8/28/97	3,300		
B12 & B9 Beaches	8/28/97	900		
Hope Lake Beach	8/14/97	300		
Peace River	8/16/97	700	1,400	0.1%
Lake Mikchalk		1,000	6,200	0.4%
Narrows				
Northshore Beaches	8/16/97	1,000		
Southshore Beaches				

(Continued)

Table 10. (Continued)

Area	Date	Aerial Count ¹	Population Estimate	Distribution %
Wind River	8/16/97	300	600	0.0%
Lake Kulik		33,200	207,100	13.7%
K1 & K2 Creeks	8/14/97	5,700		
K5 Creek - Grant River Beaches				
Grant River - K2 Creek Beaches	8/28/97	19,700 ^c		
Southshore Beaches	8/28/97	7,800		
Grant River	8/16/97	19,800	39,600	2.6%
Total		417,325	1,512,400	100.0%

- ¹ All counts rounded to the nearest 10 fish.
- ² Access blocked by beaver dams.
- ^a Ground survey counts conducted by FRI, University of Washington..
- ^b Partial count.
- ^c Combined count from K-2 through K-5 North shore.
- ^d Combined count from Hardluck bay to Peace River.

Table 11. Peak aerial counts of live sockeye salmon and total escapement estimates, Togiak District, 1997.

Stream	Aerial Counts		Total Escapement Estimate	
	Date	Number	Factor ¹	Number
<u>Togiak Section</u>				
Togiak Tower				131,682
Togiak River mainstem	19-Aug	8,200	1.5	12,300
Gechiak Lake System	22-Aug	1,600	1.5	2,400
Pungokepuk Lake	22-Aug	450	1.5	675
Nayorurun River	22-Aug	50	1.5	75
Kemuk River	19-Aug	650	1.5	975
Ongivinuk Lake System	19-Aug	2,800	1.5	4,200
Subtotal		13,750		20,625
<u>Kulukak Section</u>				
Kulukak River	23-Aug	1,700	1.5	2,550
Kulukak Lake	23-Aug	600	1.5	900
Tithe Creek Ponds	23-Aug	3,000	1.5	4,500
Subtotal		5,300		7,950
<u>Matogak, Osviak, and Cape Pierce Sections</u>				
Matogak River ²	03-Aug	360	2.0	720
Osviak River ²	03-Aug	780	2.0	1,560
Slug River ²	19-Jul	1,045	2.0	2,090
Subtotal		2,185		4,370
<u>Other</u>				
Quigmy River				
Negukthlik River ²	02-Aug	1,000	2.0	2,000
Ungalikthluk River				
Subtotal		1,000		2,000
Total		22,235		166,627

¹ Derived by expanding peak live count to reflect fish not counted due to variables such as schooled and dead fish, late or poor survey conditions, bad weather, etc..

² USFWS estimate. Sockeye salmon count obtained during chinook and chum surveys.

Table 12. Peak aerial counts of live chinook salmon and total escapement estimates, Togiak District, 1997.

River	Aerial Counts		Total Escapement Estimates	
	Date	Number	Factor ¹	Number
Togiak Section				
Togiak River mainstem				
A	28-Jul	100	2.0	200
B	28-Jul	350	2.0	700
C	28-Jul	1,300	2.0	2,600
D	28-Jul	600	2.0	1,200
E	28-Jul	820	2.0	1,640
F	28-Jul	1,000	2.0	2,000
Subtotal		4,170		8,340
Gechiak River	28-Jul	275	2.0	550
Pungokepuk River	28-Jul	180	2.0	360
Nayorurun River	28-Jul	150	2.0	300
Kemuk River	28-Jul	275	2.0	550
Ongivinuk River	28-Jul	100	2.0	200
Subtotal		980		1,960
Togiak River Drainage Total		5,150		10,300
Kulukak Section				
Kulukak River ²	28-Jul	310	2.5	775
Matogak, Osviak, and Cape Pierce Sections				
Matogak River ²	25-Jul	50	2.5	125
Osviak River ²	25-Jul	65	2.5	163
Slug River ²	31-Jul	33	2.5	83
Subtotal				370
Other				
Quigmy River ²	25-Jul	10	2.5	25
Negukthlik River				
Ungalikthluk River ²		10	2.5	25
Subtotal				50
Total				11,495

¹ Derived by expanding peak live count to reflect fish not counted due to variables such as schooled and dead fish, late or poor survey conditions, bad weather, etc..

² USFWS estimate.

Table 13. Peak aerial counts of live chum salmon and total escapement estimates, Togiak District, 1997.

River	Aerial Counts		Total Escapement Estimate	
	Date	Number	Factor ¹	Estimate
<u>Togiak Section</u>				
Togiak River mainstem				
A	28-Jul	3,900	2.0	7,800
B	28-Jul	3,100	2.0	6,200
C	28-Jul	3,800	2.0	7,600
D	28-Jul	2,750	2.0	5,500
E	28-Jul	7,100	2.0	14,200
F	28-Jul	4,550	2.0	9,100
Subtotal		25,200		50,400
Gechiak River	28-Jul	3,200	2.0	6,400
Pungokebuk River	28-Jul	800	2.0	1,600
Nayorurun River	28-Jul	4,750	2.0	9,500
Kemuk River	28-Jul	1,800	2.0	3,600
Ongivinuk River	28-Jul	3,900	2.0	7,800
Subtotal		39,650		79,300
<u>Kulukak Section</u>				
Kulukak River ²	28-Jul	4,550	2.0	9,100
<u>Matogak, Osviak, and Cape Pierce Sections</u>				
Matogak River ³	25-Jul	3,000	2.0	6,000
Osviak River ³	25-Jul	2,500	2.0	5,000
Slug River ^{2,3}	31-Jul	1,890	2.0	3,780
Subtotal		7,390		14,780
<u>Other</u>				
Quigmy River ³	25-Jul	1,700	2.0	3,400
Negukthlik River			2.0	
Ungalikthluk River			2.0	
Subtotal		1,700		3,400
Total		53,290		106,580

- 1 Derived by expanding peak live count to reflect fish not counted due to variables such as schooled and dead fish, late or poor survey conditions, bad weather, etc..
- 2 U.S. Fish and Wildlife Service estimate. Surveys were past peak of spawning, estimate is the number
- 3 Surveys were past peak of spawning; estimate is the number of live chums observed times the expansion factor, plus the number of chum carcasses counted.

Table 14. Peak aerial counts of live coho salmon and total escapement estimates, Togiak District, 1997.

Stream	Aerial Counts		Total Escapement Estimate	
	Date	Number	Factor ¹	Number
<u>Togiak Section</u>				
Togiak River mainstem				
A	29-Sep	600	3.0	1,800
B	29-Sep	200	3.0	600
C	29-Sep	400	3.0	1,200
D	29-Sep	100	3.0	300
E	29-Sep	400	3.0	1,200
F	29-Sep	1,800	3.0	5,400
Subtotal		3,500		10,500
Gechiak River	29-Sep	1,000	3.0	3,000
Pungokebuk River	29-Sep	650	3.0	1,950
Nayorurun River	29-Sep	350	3.0	1,050
Kemuk River	29-Sep	475	3.0	1,425
Ongivinuk River	29-Sep	900	3.0	2,700
Subtotal		3,375		10,125
<u>Kulukak Section</u>				
Kulukak River	30-Sep	1,675	3.0	5,025
<u>Matogak, Osviak, and Cape Pierce Sections</u>				
Matogak River ²	19-Aug	150	3.0	450
Osviak River ²	02-Oct	1,046	3.0	3,138
Slug River ²	30-Sep	1,397	3.0	4,191
Subtotal		2,593		7,779
<u>Other</u>				
Quigmy River ²	30-Sep	325	3.0	975
Negukthlik River			3.0	
Ungalikthluk River	30-Sep	1,690	3.0	5,070
Subtotal		2,015		6,045
Total		13,158		39,474

¹ Derived by expanding peak live count to reflect fish not counted due to variables such as schooled and dead fish, late or poor survey conditions, bad weather, etc..

² U.S.F.W.S. survey, includes schooled fish, indicating pro-peak timing. Negukthlik & Ungalikthluk Rivers combined.




Figure 1. Bristol Bay management area, Alaska.




Figure 2. Alagnak River drainage, Bristol Bay, Alaska.




Figure 5. Wood River Lakes system, Bristol Bay, Alaska.




Figure 6. Lake Nunavagaluk system, Bristol Bay, Alaska.




Figure 7. Nushagak-Mulchatna River system, Bristol Bay, Alaska.




Figure 8. Tikchik Lakes system, Bristol Bay, Alaska.




Figure 9. Togiak River system, Bristol Bay, Alaska.




Figure 10. Kulukak River system, Bristol Bay, Alaska.

Appendix Table 1. Sockeye salmon total escapement estimates, Naknek-Kvichak District, 1955-1997.
Estimates based on visual counts from towers unless otherwise noted.

Year	Kvichak	Naknek	Alagnak	Total	Alagnak Percent of Total
1955	250,546	278,500 ^b	171,500 ^a	700,546	24
1956	9,443,318	1,772,595 ^b	784,000 ^a	11,999,913	7
1957	2,842,810	634,645 ^b	126,595	3,604,050	4
1958	534,785	278,118	94,650	907,553	10
1959	680,000	2,231,807	825,431	3,737,238	22
1960	14,630,000	828,381	1,240,530	16,698,911	7
1961	3,705,849	351,078	90,036	4,146,963	2
1962	2,580,884	723,066	90,630	3,394,580	3
1963	338,760	905,358	203,304	1,447,422	14
1964	957,120	1,349,604	248,700	2,555,424	10
1965	24,325,926	717,798	175,020	25,218,744	1
1966	3,775,184	1,016,445	174,336	4,965,965	4
1967	3,216,208	755,640	202,626	4,174,474	5
1968	2,557,440	1,023,222	193,872	3,774,534	5
1969	8,394,204	1,331,202	122,490	9,847,896	1
1970	13,935,306	732,502	177,060	14,844,868	1
1971	2,387,392	935,754	187,302	3,510,448	5
1972	1,009,962	586,518	151,188	1,747,668	9
1973	226,554	356,676	35,280	618,510	6
1974	4,433,844	1,241,058	214,848	5,889,750	4
1975	13,140,450	2,026,686	100,480	15,267,616	1
1976	1,965,282	1,320,750	81,822	3,367,854	2
1977	1,341,144	1,085,856	100,000 ^a	2,527,000	4
1978	4,149,288	813,378	229,400 ^a	5,192,066	4
1979	11,218,434	925,362	294,200 ^a	12,437,996	2
1980	22,505,268	2,644,698	297,900 ^a	25,447,866	1
1981	1,754,358	1,796,220	82,210 ^a	3,632,788	2
1982	1,134,840	1,155,552	239,300 ^a	2,529,692	9
1983	3,569,982	888,294	96,220 ^a	4,554,496	2
1984	10,490,670	1,242,474	215,370 ^a	11,948,514	2
1985	7,211,046	1,849,938	118,030 ^a	9,179,014	1
1986	1,179,322	1,977,645	230,180 ^a	3,387,147	7
1987	6,065,880	1,061,806	154,210 ^a	7,281,896	2
1988	4,065,216	1,037,862	194,630 ^a	5,297,708	4
1989	8,317,500	1,161,984	196,760 ^a	9,676,244	2
1990	6,970,020	2,092,578	168,760 ^a	9,231,358	2
1991	4,222,788	3,578,508	277,589 ^a	8,078,885	3
1992	4,725,864	1,606,650	226,643 ^a	6,559,157	3
1993	4,025,166	1,535,658	347,975 ^a	5,908,799	6

(Continued)

Appendix Table 1. (Continued)

Year	Kvichak	Naknek	Alagnak	Total	Alagnak Percent of Total
1994	8,337,840	990,810	242,595 ^a	9,571,245	3
1995	10,038,720	1,111,140	215,713 ^a	11,365,573	2
1996	1,450,578	1,078,098	306,750	2,835,426	11
1997	1,503,732	1,025,664	218,115	2,747,511	8
Mean	5,704,988	1,239,466	241,530	7,185,984	5

^a Aerial survey counts.

^b Weir counts.

Appendix Table 2. Aerial survey counts of chinook salmon escapements, Naknek River drainage, 1970-1997.

Year	Mainstem Naknek River	Paul's Creek	King Salmon Creek	Big Creek	Total
1970	3,060		260	825	4,145
1971	1,639	52	704	490	2,885
1972	351	156	1,224	1,060	2,791
1973	1,315		115	1,106	2,536
1974		91	495	860	1,446
1975	2,250	144	279	779	3,452
1976	5,950	31	180	970	7,131
1977	4,830		1,860		6,690
1978					^a
1979					^a
1980	300	17		30	347
1981	2,890		591	790	4,271
1982	5,360	340	980	1,930	8,610
1983	2,860	290	460	4,220	7,830
1984	790	400	385	3,420	4,995
1985	590				590
1986	2,200	73	102	1,542	3,917
1987	2,800	7	290	1,353	4,450
1988	7,380	150	600	3,600	11,730
1989	1,700	50	100	860	2,710
1990	4,500	150	350	2,000	7,000
1991	1,655	121	275	2,340	4,391
1992	1,550	88	158	895	2,691
1993	5,520	86	700	1,710	8,016
1994	5,970	203	974	2,531	9,678
1995	2,790	26	239	1,905	4,960
1996	2,965	157	312	1,576	5,010
1997	7,520	248	902	1,783	10,453
Mean	3,281	120	522	1,607	5,530 ^b
Percent	59	2	9	29	100

^a Counts unavailable.

^b The sum of mean indices.

Appendix Table 3. Chinook salmon escapement survey history, mainstem Naknek River, 1929-1997.

Year	Count Dates	Surveyors	Actual Weir Count ¹	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ²	Comments
1929	7/03-7/31		1,498			Chinook count peaked 7/27.
1930	6/20-8/09		1,999			Chinook count peaked 8/09.
1931	6/17-8/09		896			Chinook count peaked 8/07.
1932	6/23-8/10		1,869			
1950	7/08-8/20		3,097			Chinook count peaked 8/09.
1951	6/28-8/07		1,876			Chinook count peaked 8/04.
1952	6/25-8/10		633			Chinook count peaked 8/06.
1953	6/24-8/10		2,074			Chinook count peaked 7/26.
1954	6/20-8/11		3,474			Chinook count peaked 8/10.
1955	6/13-8/17		4,188			Chinook count peaked 8/16.
1956	6/22-8/28		7,378			Chinook count peaked 8/18.
1957	6/28-8/04		8,504			Chinook count peaked 8/03.
1966		Redick				300 were counted 8/26 from a skiff in the Rapids.
1967	Mid-Aug.	Paddock			800	
1968					1,200	Conservative estimate.
1969					1,200	
1970	7/31	Whitehead		845		
	8/03	Siedelman		3,060		Visibility very good. Super-cub.
	8/22	Siedelman		1,540	1,750	Water high & murky. Spawning pre-peak.
	8/22	Whitehead		1,310		
	8/25	Whitehead		2,225		Counting conditions optimal.
	8/25	Siedelman		2,536	2,500	Conditions good. Spawning pre-peak.

(Continued)

Appendix Table 3. (Continued)

Year	Count Dates	Surveyors	Actual Weir Count ¹	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ²	Comments
1971	8/26	Cunningham		1,639		Fish concentrated near Rapids Camp. Few dead.
1972	8/23	Cunningham & McCurdy		351		Poor counting conditions. Post-peak.
1973	8/19	Russell		1,315		Counting conditions good. Peak near at hand.
1974	8/19	Russell			450	Count accuracy questionable. Many fish were deep.
1975	8/17	Russell		2,250		Good viewing, peak near. Still fish spawning 9/08.
1976	8/13	Bill		2,615		Spawning near peak. Very few dead.
	8/16	Russell		5,950	7,250	Pre-peak. Still lots fish holding in large groups.
1977	8/22	Russell		4,830	5,750	Pre-peak. Few dead. Some still holding deep.
1978	8/09	Gwartney			4,000	Near peak.
1983	8/14	Bill		2,860	3,000	Pre-peak. Still fish holding in large groups.
1984	8/14	Bill		790	2,370	
1985	8/06	Bill			600	Pre-peak.
	8/27	Bill		590	700	
1986	8/18	Russell		1,990		Spawning pre-peak. Still many fish holding.
	8/19	Meyer		2,200		Peak of spawning drawing near.
1987	8/19	Meyer		2,800		Pre-peak. Fish still in large groups. Few redds.
	8/28	Bill		2,655	2,855	
1988	8/09	Minard		7,380	7,400	Approaching peak. Most fish on redds.
1989	8/14	Minard		1,700		Fish actively spawning. Few carcasses observed.
1990	8/06	Minard		4,500		
1991	8/20	Russell		1,655		Pre-peak. Still many fish schooled & waiting.

(Continued)

Appendix Table 3. (Continued)

Year	Count Dates	Surveyors	Actual Weir Count ¹	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ²	Comments
1992	8/21	Regnart		877		Water clarity poor in deeper pools.
	8/27	Regnart		1,550		At Peak...all fish on redds.
1993	8/23	Regnart		5,520		Near peak. Still some fish schooled.
1994	8/24	Regnart		5,970		Near peak. Most on redds.
1995	8/21	Regnart		2,790		Near peak. Most on redds.
1996	8/21	Regnart		2,965		At Peak...all fish on redds.
1997	8/16	Regnart		7,520		Near peak. Most on redds.
Mean			3,124	2,799		

¹ Weir count did not account for estimated 15-20% of chinook that spawn downstream of weir site. Also does not account for fish that migrated upstream past the weir site before and after weir operation.

² Surveyor's subjective estimate of instantaneous population of chinook salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting at time of survey.

Appendix Table 4. Chinook salmon escapement survey history, Big Creek, Naknek River Drainage, 1963-1997.

Year	Count Dates	Surveyors	Float Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ¹	Comments
1963	8/01	Paddock		362		Covered only half stream length. Helicopter.
	8/13	Paddock		1,345	2,690	Spawning near peak. Good survey.
1964	7/31	Paddock		484		Survey too early.
	8/15	Siedelman & Williamson		636		Survey fair to good. Near peak. Helicopter.
	8/15-8/18	Siedelman & Williamson	1,130			Peak of spawning over.
1965	8/05-8/08	Andrews	578			Fair survey. Began below Index Area No. 1.
1966	8/13-8/16	Redick	979			Spawning at peak. Included Index Area No. 1. Count affected by rain/turbid water in lower areas.
1967	8/10-8/14	Whitehead & Bury	1,129			Upstream redds occupied while those in the lower stream area were abandoned.
1968	8/10-8/14	Meyers & Preyer	3,827			Counting conditions fair to poor.
1969	8/12-8/14	Parkinson & Faro	1,012			High murky waters hampered float count.
	Mid-Aug.	??			5,000	Flown due to poor count conditions during float.
1970	7/19	Whitehead		825		
	8/15-8/17	Parkinson & Brooks	1,601			High murky waters in lower 2/3 of stream.

(Continued)

Appendix Table 4. (Continued)

Year	Count Dates	Surveyors	Float Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ¹	Comments
1971	8/13	Cunningham		490	1,200	Only upper 1/3 of stream surveyed due to murky water in lower 2/3.
	8/28	Siedelman		277		Past peak. Survey affected by winds of 30+ mph.
1972	8/08	Cunningham		695		Pre-peak.
	8/18	Siedelman		1,060		Post-peak.
1973	8/17	Russell		1,106		At peak of spawning. Many fish beaten up (fungus).
1974	8/01	Russell		520	850	Pre-peak. No dead chinook. Lots dead chums.
	8/11	Russell		860	1,250	Didn't survey lower 8 miles of creek 8/11. Could add 150 fish to survey as Russell saw that many in the unsurveyed portion from skiff 8/10. Near peak.
1975	8/09	Russell		779		Survey pre-peak.
1976	8/13	Bill		970	1,400	Not total stream coverage due to winds & low fuel.
1983	8/14	Bill		4,220	9,000	
1984	8/08	Bill		3,420	8,800	At peak of spawning.
1985	8/06	Bill			2,900	Survey conditions..high water & gusty winds.
1986	8/08	Meyer		1,542	6,000	Excellent conditions. Fish at spawning peak.
1987	8/21	Meyer		1,353	2,500	
1988	8/09	Minard		3,600		
1989	8/14	Minard		860		
1990	8/06	Minard		2,000		
1991	8/12	Regnart		2,340		At spawning peak..all fish on redds, only 20 dead.
1992	8/18	Regnart		895		Est. 5-6 days post-peak. Count includes 125 dead.

(Continued)

Appendix Table 4. (Continued)

Year	Count Dates	Surveyors	Float Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ¹	Comments
1993	8/17	Regnart		1,710		Estimated survey 3-4 days past peak.
1994	8/16	Regnart		2,531		Est. 2-3 days post-peak. Count includes 159 dead.
1995	8/15	Regnart		1,905		Estimate survey was several days past peak.
1996	8/12	Regnart		1,576		At spawning peak....38 dead observed
1997	8/7	Regnart		1,783		At spawning peak....48 dead observed
Mean			1,465	1,384		

¹ Surveyor's subjective estimate of instantaneous population of chinook salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting at time of survey.

Appendix Table 5. Chinook salmon escapement survey history, King Salmon Creek, Naknek River drainage, 1964-1997.

Year	Count Dates	Surveyors	Float Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ¹	Comments
1964	7/31	Paddock		378		Survey conditions fair. Helicopter.
	8/11	Paddock		55		Visibility poor. Helicopter.
	8/11-8/14	Paddock & Siedelman	104			Peak of spawning long past. Poor survey (turbid).
1966	7/31-8/03	Redick	633			Spawning at or near peak.
1967	7/24-7/26	Paddock	289			Poor visibility. Estimated 600 fish present.
1968	7/17	Whitehead		282		Pre-peak. Helicopter.
	7/17	Meyers		242		Pre-peak. Helicopter.
	7/20	Whitehead		868		Optimum coditions. Count from H-21 Helicopter.
	7/20	Meyers		575		Optimum coditions. Count from H-21 Helicopter.
	7/20-7/23	Whitehead & Meyers	2,204			Counting conditions optimum.
1969	7/23-7/25	Parkinson & Berry	2,722			Pre-peak. Count fair-to-poor last 2 days (weather).
1970	7/19	Whitehead		260		Counting conditions poor. Pre-peak.
1971	7/28	Cunningham		704		Visibility was good.
1972	7/29	Siedelman		1,224		Peak of spawning.
1973	8/01	Siedelman		115		Visibility only fair. Survey possibly post-peak.
1974	7/15	Russell		164	350	Pre-peak. Many fish holding in pools.
	7/28	Russell		495	625	At or near peak. Only one carcass obsd. Good vis.
1975	7/28	Russell		279	375	Survey pre-peak. Good viewing conditions.
	8/10	Russell	67			Floated only lower 12 miles of creek.
	8/17	Russell		0		Excellent viewing conditions. Spawning is done.
1976	8/03	Bill		180	400	Peak within next 3 days.
1977	7/29	Russell		1,860	2,350	At peak of spawning.

(Continued)

Appendix Table 5. (Continued)

Year	Count Dates	Surveyors	Float Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ¹	Comments
1978	8/09	Gwartney			350	Past peak. Viewing good. Most fish dead or spent.
1979	??	Gwartney			1,750	
1980	8/08	Bill				Creek too high & muddy to census.
1981	7/30	Russell		591	1,500	Peak of spawning in progress. Vis = fair-to-poor.
1982	8/07	Bill		980	3,920	Good visibility.
1983	8/14	Bill		460	1,400	Poor visibility. Muddy. 30% spawners dead already.
1984	8/08	Bill		385	1,155	
1988	8/08	Minard		600		At peak.
1989	8/14	Minard		100		Past peak.
1990	8/06	Minard		350		
1991	7/30	Russell		100		Pre-peak and water clarity only "Fair".
	8/05	Russell		275		Est. at spawning peak, most fish on redds, 2 dead.
1992	8/09	Russell		158		Post-peak as 47 dead counted & aband. redds numerous.
1993	7/31	Russell		700	900	Slightly pre-peak. Most fish on redds. Water clear.
1994	7/29	Russell		974		Slightly pre-peak. Most fish on redds. Only 6 carcasses.
1995	8/05	Russell		239		A little past peak. Several singles on redds. Vis. only
1996	8/05	Regnart		312		Slightly post peak. 26 dead counted.
1997	7/18	Regnart		902		Pre-peak and water clarity "Good".
Mean			1,190	478		

¹ Surveyor's subjective estimate of instantaneous population of chinook salmon spawners in the river at time of aerial survey, based on survey coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for conditions, river area counting at time of survey.

Appendix Table 6. Chinook salmon escapement survey history, Paul's Creek, Naknek River drainage, 1971-1997.

Year	Count Dates	Surveyors	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ¹	Comments
1971	7/28	Cunningham	52		
1972	7/28	Siedelman	156		Prior to peak.
1973	8/01	Siedelman			Too murky to survey.
1974	7/15	Russell	2		
	7/26	Russell	91	250	Prior to spawning peak.
1975	7/28	Russell	144	225	Prior to peak. Good conditions.
1976	8/03	Bill	31	100	Poor conditions. Fish paired & spawning.
1977					No count.
1978	8/09	Gwartney		300	Past peak. 75% of fish dead.
1979					No count.
1980	8/08	Bill	17		All carcasses. Creek high & muddy.
1981					No count.
1982	8/07	Bill	340	1,020	Good visibility. Spawning near peak.
1983	8/14	Bill	290	800	Poor visibility.
1984	8/08	Bill	400	800	Fair visibility. About 25 % dead already.
1985	8/06	Bill		170	Pre-peak.
1986	8/08	Meyer	73	236	Approximately 30 % dead already.
1987	8/13	Russell	7		Poor survey conditions. Past peak.
	??	Meyer		400	Estimat 400 present based on jet boat surveys.
1988	8/08	Minard	150		At peak.
1989	8/14	Minard	50		Past peak. Excellent visibility.
1990	8/06	Minard	150		Excellent survey conditions.

(Continued)

Appendix Table 6. (Continued)

Year	Count Dates	Surveyors	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ¹	Comments
1991	7/30	Russell	121		Slightly pre-peak. Only 1 carcass noted.
1992	8/01	Russell	88		Slightly pre-peak. Stream clarity only "Fair".
1993	7/31	Russell	86	140	Slightly pre-peak. Overflow approx 60% of stream.
1994	7/29	Russell	203	300	Pre-peak...but many fish on redds.
1995	8/05	Russell	26		Water clarity poor. 5 carcasses noted
1996	8/05	Regnart	157		Peak of spawning. 12 dead counted.
1997	7/18	Regnart	248		Pre-peak. Excellent visibility
Mean			131		

¹ Surveyor's subjective estimate of instantaneous population of chinook salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting at time of survey.

Appendix Table 7. Chinook salmon escapement survey history, Alagnak River, 1963-1997.

Year	Count Dates	Surveyors	Float Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ¹	Comments
1963	8/12	Siedelman		551		Excellent conditions. No side channels flown.
1966	8/06	Redick		13		Poor conditions.
	8/06-8/10	Redick	238			Nonvianuk & mainstem portions only (not Kukaklek).
	8/11	Redick		1,465		Pre-peak. Still many fish upmigrating.
1967	8/16	VanValin		1,250		
1968	8/18	Siedelman		6,717	8,500	Fairly good survey.
1969	8/19	Siedelman		4,781	6,000	Marginal survey conditions, (20kn NW winds).
1970	8/22	Siedelman		5,250	5,000	Peak of spawning. Visibility good
	8/22	Whitehead		4,590		Peak of spawning. Visibility good
1971	8/25	Siedelman		1,420	1,500	Water high, but count okay.
	8/25	Cunningham		1,475		
1972	8/23	Cunningham		2,256	2,400	Past peak. Many dead. Many unoccupied redds.
1973	8/16	Russell		824	1,250	Near peak of spawning. No dead though.
1974	8/13	Russell		1,411	1,700	Pre-peak.
	8/19	Russell		1,596	1,900	Spawning near peak.
1975	8/17	Russell		6,620	7,250	About a week pre-peak. Some large groups holding.
1976	8/16	Bill		7,593	8,750	Pre-peak. Not many dead yet.
1977	8/18	Bill		3,634	12,000	Pre-peak. Didn't count river below Pfaff Pond.
	8/18	Sanders		9,425		Pre-peak. Didn't count river below Pfaff Pond.
1978	8/24	Bill		11,650	25,100	
1979						No survey.
1980	8/08	Bill		2,020	5,090	Pre-peak. Fog over lower river.
	8/21	Bill		2,930	5,860	
1981	8/26	Bill		2,430	8,540	

(Continued)

Appendix Table 7. (Continued)

Year	Count Dates	Surveyors	Float Count	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ¹	Comments
1982	8/09	Bill		3,400	4,700	At least a weak too early.
	8/19	Bill		3,350	5,480	Peak survey.
1983	8/15	Bill		2,980	3,500	At peak of spawning.
1984	8/14	Bill		6,090	9,135	
1985	8/17	Bill		3,920	9,518	About peak for chinook spawning. 30% dead already.
1986	8/11	Bill		3,090	7,200	Peak of spawning.
1987	8/22	Bill		2,420		
1988	8/12	Bill		4,600		
1989	8/15	Bill		3,650		
1990	8/08	Bill		1,720		
1991	8/09	Regnart		2,023		Pre-peak. Most fish schooled yet. Few on redds.
	8/19	Regnart		2,531		Near peak. Most fish on redds.
1992	8/10	Regnart		3,042		Pre-peak. Most fish still schooled.
	8/21	Regnart		2,275		Near peak...but water clarity worse than earlier.
1993	8/09	Regnart		10,170		Near peak. Most on redds.
1994	8/08	Regnart		8,480		About half the fish on redds. Others schooled.
1995	8/10	Regnart		6,860		About 2/3 of chinook noted on redds.
1996	8/12	Regnart		9,885		Near peak. Most on redds.
1997	8/7	Regnart		15,210		Peak. Excellent visibility
Mean			238	4,283		

¹ Surveyor's subjective estimate of instantaneous population of chinook salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting at time of survey.

Appendix Table 8. Chinook salmon escapement survey history, Kvichak River, 1932-1997.

Year	Count Dates	Surveyors	Weir Count	Non-expanded Aerial Index Count	Expanded Aerial Inde Estimate ¹	Comments
1932	6/28-8/5		5,753			Peak count was on 7/05 (1,168 fish).
1976	8/16	Bill		35	45	Survey timed to count pink salmon.
1980 ^a	8/08	Bill		900	1,000	Chinook actively spawning.
1984	8/14	Bill		200		
1988	8/13	Bill		190	570	Nearly all on redds.
1989	8/16	Bill		100	260	
1990	8/19	Bill		170	510	
1992	8/13	Regnart		264		All fish on redds in Kaskanak Flats.
1993	8/16	Regnart		115		All fish on redds in Kaskanak Flats.
1994	8/12	Regnart		306		
1995	8/14	Regnart		96		
1996	8/18	Regnart		132		
1997	8/15	Regnart		103		
Mean			5,753	218		

¹ Surveyor's subjective estimate of instantaneous population of chinook salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting at time of survey.

^a Pecks Creek, a Kvichak River tributary, was float surveyed 7/30-8/03, 1980 by R. Russell and 99 spawning chinook salmon were counted.

Appendix Table 9. Chinook salmon escapement data, Naknek-Kvichak District, 1970-1997.

Year	Non-expanded Escapement Indices by Drainage ¹			Total
	Naknek	Alagnak	Kvichak	
1970	4,145 ^a	5,250		9,395
1971	2,885	1,420		4,305
1972	2,791	2,256		5,047
1973	2,536 ^a	824		3,360
1974	1,446 ^b	1,596		3,042
1975	3,452	6,620		10,072
1976	7,131	7,593	35	14,759
1977	6,690 ^a	3,634		10,324
1978	^c	11,650		11,650
1979	^c			0
1980	347 ^d	2,930	900	4,177
1981	4,271 ^a	2,430		6,701
1982	8,610	3,400		12,010
1983	7,830	2,980		10,810
1984	4,995	6,090	200	11,285
1985	590 ^e	3,920		4,510
1986	3,917	3,090		7,007
1987	4,450	2,420		6,870
1988	11,730	4,600	190	16,520
1989	2,710	3,650	100	6,460
1990	7,000	1,720	170	8,890
1991	4,391	2,531		6,922
1992	2,691	3,042	264	5,997
1993	8,016	10,170	115	18,301
1994	9,678	8,480	306	18,464
1995	4,960	6,860	96	11,916
1996	5,010	9,885	132	15,027
1997	10,453	15,210	103	25,766
Mean	5,105	5,164	100	10,369

¹ Includes aerial indices from all streams surveyed in drainage.

^a No index count for Paul's Creek.

^b No index count for Naknek River.

^c No non-expanded index counts exist for this year.

^d Includes only index counts for mainstem Naknek River, Paul's Creek, & Big Creek.

^e Naknek River mainstem only.

^f Sum of mean indices.

Appendix Table 10. Chum salmon escapement survey history, Alagnak River, 1961-1997.

Year	Count Dates	Surveyors	Tower Counts	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ¹	Comments
1961			18,906			
1962			3,846			
1963	8/12	Siedelman	20,124	4,120		
1964			2,562			
1965			132			
1966						
1967			9,990			
1968			72			
1969			210			
1970			5,790			
1971			402			
1972			48			
1973						
1974						
1975						
1976	8/16	Bill		2,125	5,250	
1977	8/18	Bill		35,000		
1978	8/24	Bill		9,900		
1979						
1980	8/21	Bill		7,300	14,600	
1981	8/26	Bill		75,000	75,000	
1982	8/09	Bill		14,000	42,000	
	8/19	Bill		12,000	30,000	
1983	8/15	Bill		8,800		Pre-peak.

(Continued)

Appendix Table 10. (Continued)

Year	Count Dates	Surveyors	Tower Counts	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ¹	Comments
1984	8/14	Bill		48,000	87,500	
1990	8/08	Bill		8,500	30,000	Pre-peak.
	8/18	Bill		48,800		Close to peak of spawning.
1991	8/09	Regnart		43,000		Pre-peak.
	8/19	Regnart		64,300		Peak of spawning.
1992	8/10	Regnart		114,000		Near Peak.
1993	8/09	Regnart		4,600		Near Peak.
1994	8/08	Regnart		62,900		Near Peak.
1995	8/10	Regnart		132,000		Near Peak.
1996	8/12	Regnart		145,000		Near Peak
1997	8/7	Regnart		37,800		Near Peak
Mean			3,575	43,857		

¹ Surveyor's subjective estimate of instantaneous population of chum salmon spawners in the river at time of aerial survey, based on survey conditions, coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for river area counting at time of survey.

Appendix Table 11. Pink salmon escapement survey history, Alagnak River, 1968-1997.

Year	Count Date	Surveyor	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ¹	Comments
1968	8/27	Siedelman	97,000	125,000	
1970					No survey.
1972					No survey.
1974	8/14	Bill	20,600		Big schools. Pre-peak.
1976	8/16	Bill	6,375	13,000	Pre-peak.
1978	8/24	Bill	330,300	736,000	Just starting to spawn. Many still in lower river.
1980	8/21	Bill	121,000	242,000	
1982	8/09	Bill	21,300	63,900	
	8/19	Bill	24,800	43,000	Pre-peak.
1984	8/14	Bill	296,500	567,100	Survey too early for peak. Most fish schooled.
1986	8/11	Bill	48,600	145,800	
1988	8/12	Bill	415,000	620,000	Pre-peak.
1990	8/08	Bill	45,100		
	8/18	Bill	240,500		Estimated to be about 1 week pre-peak.
1992	8/10	Regnart	15,000		Pre-peak.
1993	8/09	Regnart			No pinks noted.
1994	8/08	Regnart			No pinks noted.
1995	8/10	Regnart			No pinks noted.
1996	8/12	Regnart			No pinks noted.
1997	8/7	Regnart			No pinks noted.
Mean			146,880		

¹ Surveyor's subjective estimate of instantaneous population of pink salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting a time of survey.

Appendix Table 12. Pink salmon escapement survey history, Kvichak River, 1966-1997.

Year	Count Dates	Surveyor	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ¹	Comments
1966		Robertson		67,500	
1968	8/26	Siedelman		88,000	
1970					No survey.
1972					No survey.
1974	8/14	Bill		30,560	
1976	8/16	Bill		16,100	Most still schooled.
1978	8/28	Bill	88,000	440,000	Still numerous fish migrating & some schooled.
1980	8/08	Bill	7,000	25,000	Still schooled.
1982					No Survey.
1984	8/14	Bill	111,000	165,000	
1986					No survey.
1988	8/13	Bill	94,000		
1990	8/19	Bill	25,300	47,000	
1992					No survey.
1993					No survey.
1994					No survey.
1995					No survey.
1996					No Survey
1997					No Survey.
Mean			65,060		

¹ Surveyor's subjective estimate of instantaneous population of pink salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting at time of survey.

Appendix Table 13. Pink salmon escapement survey history, Naknek River, 1974-1997.

Year	Count Dates	Surveyor	Non-expanded Aerial Index Count	Expanded Aerial Index Estimate ¹	Comments
1974	8/14	Bill	161,800	362,000	
1976	8/13	Bill	94,600	110,000	Just pre-peak. Many still schooled.
1978	8/24	Bill	312,000	780,000	
1980	8/08	Bill	80,000	160,000	Pre-peak.
1982	8/19	Bill	33,600	34,000	Pre-peak.
1984	8/14	Bill	27,000	125,000	
1986	8/18	Russell	286,000	375,000	Most fish still schooled and holding. Pre-peak.
1988	8/24	Russell	187,000		
1990	8/18	Bill		65,000	
1992					No survey.
1993					No survey.
1994					No survey.
1995					No survey.
1996					No Survey.
1997					No survey.
Mean			147,750		

¹ Surveyor's subjective estimate of instantaneous population of pink salmon spawners in the river at time of aerial survey, based on survey conditions, river area coverage, water clarity, etc. Does not include expansion for earlier or later run fish not available for counting at time of survey.

Appendix Table 14. Aerial survey counts of chinook salmon escapement, Egegik District, 1981-1997. ^a

Year	Egegik River	Shosky Creek	Whale Mountain Creek	Mossy Creek	Mink Creek	Gertrude Creek	Kaye's Creek	Takayoto Creek	Angle Creek	Contact Creek	King Salmon River	Total
1981						515						515
1982	300					900				300		1,500
1983						860		380		375		1,615
1984	40	300				600		350		110		1,400
1985	75	80	0	15	10	260	230	315		95		1,080
1986	65	150	48	0	0	150	46	40		18	15	532 ^b
1987	15	174	2	74	0	408	284	232	2	88		1,279
1988	50	151	0	12		248	120	177		110		868
1989	14	90	13	43	7	310	120	300		100		997
1990	24 ^c	85	7	35	2	260	175	175		205		968
1991	0 ^c	62	60	30	33	83	117	95		73		553
1992 ^d	15	143	52	54	22	416	320	190		296		1,508
1993	80	58	6	38	6	350	170	200		235		1,143
1994 ^d	66 ^c	48	32	118	77	840	214	230		705		2,330
1995 ^d	60 ^c	32	10	53	103	456	248	130		275		1,367
1996	42 ^c	102	8	38	20	230	74	123	6	203		846
Average	60	113	20	43	25	430	177	210	4	213		1,156
1997	30 ^c	39	2	18	10	260	173	374		740		1,646

^a Peak aerial counts unless otherwise noted. Data not expanded.

^b Survey 10-14 days later than normal.

^c Tower count.

^d Helicopter surveys.

Appendix Table 15. Aerial survey counts of chum salmon escapement, Egegik District, 1982-1997. ^a

Year	Egegik River	Shosky Creek	Whale Mountain Creek	Mossy Creek	Mink Creek	Gertrude Creek	Kaye's Creek	Takayoto Creek	Angle Creek	Contact Creek	King Salmon River	Total
1982						12,000				2,000		14,000
1983	6 ^b					5,000		3,500		6,000		14,500
1984	800	200				13,000		2,400		10,000		26,400
1985	400	0	600	200	35	2,600	800	0		500	50	5,185
1986	0	0	6,025			140	3	5	0	15	25	6,213 ^c
1987	150	0	19,000	16	1,000	3,770	2,780	0		2,850		29,566
1988	500	50	4,400	100	50	5,200	1,600	0		3,200		15,100
1989	0	10	3,200	25	100	1,100	0	0		200	14	4,649
1990	72 ^b	0	2,000	0	150	1,675	80	0		750		4,727
1991	0 ^b	0	1,500	70	100	990	280	0		480		3,420
1992 ^d	50	0	680	15	25	4,500	400	0		3,630	200	9,500
1993	100	0	1,020	8	1	1,075	0	0		100		2,304
1994 ^d	42 ^b	0	1,700	5	7	760	175	30		260		2,979
1995 ^d	144 ^b	2	395	15	30	560	162	5		600		1,913
1996	12	0	438	4	20	530	0	24	0	633		1,661
Average	175	20	3,413	42	138	3,527	523	426	0	2,081	72	9,474
1997	0 ^b	0	220	8	10	495	290	60		640		1,723

63

^a Peak aerial counts unless otherwise noted. Data not expanded.

^b Tower count.

^c Survey 10-14 days later than normal.

^d Helicopter surveys.

Appendix Table 16. Aerial survey counts of pink salmon escapement, Egegik District, 1974-1997.^a

Year	Egegik River	Whale Mountain Creek	Gertrude Creek	Contact Creek	Takayoto Creek	Kaye's Creek	Other	Total
1974	3,912 ^b							3,912
1976	0 ^b							0
1977	84 ^b							84
1980	0 ^b							0
1982	15,000							15,000
1983			58 ^c					58
1984	17,000							17,000
1985								
1986	2,500							2,500
1987								
1988	23,000							23,000
1989	300							300
1990	17,000		40 ^c					17,040
1991		88 ^d	24 ^d	36 ^d				148
1992 ^e	6 ^b	10					3	13
1993	50							50
1994	21,282 ^b							21,282
1995	24 ^b							24
1996	103,116 ^b							103,116
Average	16,937	49	41	36			3	11,972
1997	0 ^b							0

^a Non-expanded aerial peak counts unless otherwise noted.

^b Tower counts.

^c Float count.

^d Foot survey (USFWS).

^e Helicopter surveys.

Appendix Table 17. Aerial survey counts of coho salmon escapement, Egegik District, 1981-1997.

Year	Number of Surveys	Coho Salmon Count	Comments
1981	1 ^a	4,000	Only Becharof tributaries surveyed.
1982	1	20,000	Surveyed on August 20.
1983	0	0	No surveys done.
1984	3	43,225	40,000 counted in Egegik Lagoon on August 15.
1985	3	5,260	Peak surveys on August 26.
1986	1	12,575	Surveyed August 19.
1987	6	6,930	Included King Salmon River & tributaries.
1988	6	13,715	Included King Salmon River & tributaries.
1989	9	4,485	Included Gertrude & Whale Mountain Creeks.
1990	7	13,400	Peak survey on August 17.
1991	0	220	Incidental observation made August 6.
1992 ^b	0	200	Incidental observation in Egegik River August 6.
1993	0	1,130	Incidental observation from Egegik River August 16.
1994 ^{b,c}	2	7,412	Included King Salmon River & tributaries.
1995 ^d	2	5,258	Included King Salmon River & tributaries.
1996 ^e	2	9,043	Included King Salmon River & tributaries.
1997	3	4,106	Includes Gertrude Weir Count and Becharof tributaries.

^a Survey done by USFWS personnel.

^b Helicopter surveys.

^c The Egegik River Tower was maintained through September 11 and approximately 10,140 coho salmon were counted.

^d The Egegik River Tower was maintained through August 30 and approximately 7,470 coho salmon were counted.

^e The Egegik River Tower was maintained August 7 to September 11 and approximately 24,918 coho salmon were counted.

Appendix Table 18. Aerial survey counts of chinook salmon escapement, Ugashik District, 1980-1997.

Year	Ugashik River	Dog ¹ Salmon River	King Salmon	Painter Creed	Pumice Creek	Old Creek	Total
1980	0 ^a		900	1,000			1,900
1981	18 ^a		50	300			368
1982	0 ^a		700	700			1,400
1983	50 ^a	1,635	525	635	1,800	660	5,305
1984	108 ^a	836	4,100	1,875	1,100	880	8,899
1985	150 ^b	560	4,600	410	930	410	7,060
1986	66 ^b	252	1,777	646	705	739	4,185
1987	54 ^a	751	981	1,051	1,602	1,155	5,594
1988	249 ^c	900	5,820	1,170	1,025	660	9,824
1989	226 ^{bc}	848	1,670	1,030	510	520	4,804
1990	67 ^{ac}	540	1,500	590	450	610	3,757
1991	131 ^{ac}	449	700	365	375	420	2,440
1992 ^d	260 ^{ac}	821	1,260	855	750	815	4,761
1993	188 ^{ac}	579	1,970	865	450	635	4,687
1994 ^d	233 ^{ac}	1,741	2,225	1,005	2,530	1,490	9,224
1995	149 ^{ac}	882	440	366	501	505	2,843
1996	76 ^{ac}	1,079	1,200	403	^f	30 ^f	2,788
Average	119	848	1,789	780	979	681	4,696
1997	839 ^{ac}	906	802	525	536	558	4,166
Deviation ^e	604 %	7 %	-55 %	-33 %	-45 %	-18 %	-11 %

¹ Includes Figure-Eight, Goblet, Oldham, and Wandering Creeks.

^a Tower counts

^b Tower count plus later aerial survey counts of main river.

^c Survey included Grassy Creek (tributary downstream of Ugashik Lagoon).

^d Helicopter surveys.

^e 1997 deviation from 1980-1996 average.

^f Water was too turbid to see fish.

Appendix Table 19. Aerial survey counts of chum salmon escapement, Ugashik District, 1980-1997.

Year	Ugashik River	Dog ¹ Salmon River	King Salmon River	Painter Creek	Pumice Creek	Old Creek	Other	Total
1980	18 ^a		7,000	3,000				10,018
1981	0 ^a		200					200
1982	12 ^a		19,000	35,000			650	54,662
1983	0 ^a	1,650	2,700	4,000	20,000 ^b	3,300		31,650
1984	132 ^a	750	119,000	16,000	16,000	14,500	2,500	168,882
1985	42 ^c	350	20,000	1,925	6,000	670	300	29,287
1986	0 ^c	120	8,650	1,200	2,000	630	125	12,725
1987	130 ^c	340	9,750	2,290	10,340	2,090	40	24,980
1988	752 ^{c,d}	2,290	25,000	10,500	11,650	5,800	950	56,942
1989	600 ^{c,d}	1,005	7,500	3,700	2,200	2,010	625	17,640
1990	312 ^{c,d}	170	6,200	1,150	1,630	410	10	9,882
1991	315 ^{c,d}	240	7,400	750	2,550	2,525	130	13,910
1992 ^e	510 ^{a,c,d}	1,210	8,525	4,000	14,000	15,000	0	43,245
1993	93 ^{c,d}	105	7,000	720	2,040	1,025	8	10,991
1994 ^e	66 ^{a,c}	851	9,150	1,625	12,750	6,975	150	31,567
1995	6 ^{a,c}	160	3,900	1,370	2,600	1,800	0	9,836
1996	138 ^a	85	16,500	700	7,400	2,500	0	27,323
Average	184	666	16,322	5,496	7,940	4,231	392	32,573 ^f
1997	100	450	10,500	4,200	5,300	9,480	115	30,145
Deviation ^g	-46%	-32%	-36%	-24%	-33%	124%	-71%	-7%

¹ Includes Figure-Eight, Goblet, Oldham, and Wandering Creeks.

^a Tower counts

^b Float count done from a raft.

^c Survey included Grassy Creek (tributary downstream of Ugashik Lagoon).

^d Included tower count plus later aerial survey count.

^e Helicopter surveys.

^f Average of the sums of indices for all locations.

^g 1997 deviation from 1980-1996 average.

Appendix Table 20. Aerial survey counts of pink salmon escapement, Ugashik District, 1980-1997.

Year	Number of Surveys ^a	Pink Salmon Count	Comments
1980	1	2,000	
1982	1	6,000	4,000 in King Salmon River, 2,000 in Painter Creek.
1983	2	803	Survey of Dog Salmon River conducted by USFWS.
1984	3	656	650 counted in King Salmon River during September 21
1985	3	0	
1986	1	350	Observed in King Salmon River on August 19.
1987	2	1	
1988	7	2,800	Peak count on August 23: 2,000 in King Salmon River.
1989	8	50	Observed in Ugashik River on August 9.
1990	5	2,000	Peak count on August 13.
1991	0	660	Ugashik River tower count.
1992 ^b	0	1,728	Ugashik River tower count.
1994 ^b	0	425	Observed near Ugashik Lake Outlet on August 11.
1995	0	36	Ugashik River tower count.
1996	0	550	Observed in King Salmon River on August 12.
1997	0	0	

^a Zero indicates no surveys designated to look for pink salmon and any observations recorded would be incidental to surveying for other species.

^b Helicopter survey.

Appendix Table 21. Aerial survey counts of coho salmon escapement, Ugashik District, 1981-1997.

Year	Number of Surveys	Coho Salmon Counts	Comments
1981	1	13,300	Surveyed on September 7.
1982	1	10,000	Surveyed on August 26.
1983	0		
1984	1	6,100	Surveyed on August 31.
1985	2	18,880	16,500 in King Salmon River on September 12.
1986	2	8,455	Surveyed on August 19 and 25.
1987	2	17,000	16,700 in King Salmon River on August 23.
1988	7	28,280	12,900 in King Salmon River on September 7.
1989	4	11,515	7,615 observed on August 14.
1990	5	12,610	
1991	0	400	Incidental observation made August 12.
1992 ^a	0	790	Incidental observation made August 11.
1993	0	705	Incidental observation made August 16.
1994 ^a	0	760	Incidental observation made August 11.
1995	0		
1996	1	8,275	Surveyed on September 27 and 28.
1997	2	9,400	Surveyed on September 30 and October 17.

^a Helicopter survey.

Appendix Table 22. Spawner distribution and total escapement estimates of sockeye salmon, Wood River system, 1959-1997.

Year	Spawner Distribution (%)			Total Escapement ¹
	Creeks	Beaches	Rivers	
1959	32.8	50.3	16.9	2,209,300
1960	27.4	55.5	17.1	1,016,100
1961	11.4	32.3	56.3	460,700
1962	24.0	65.2	10.8	873,900
1963	12.1	68.5	19.4	721,400
1964	18.9	64.0	17.1	1,076,100
1965	40.6	11.1	48.3	675,100
1966	16.4	54.9	28.7	1,208,700
1967	9.3	66.2	24.5	515,800
1968	9.9	50.8	39.3	649,300
1969	8.6	42.4	49.0	604,300
1970	14.0	52.4	33.6	1,162,000
1971	11.2	56.8	32.0	851,200
1972	17.4	45.1	37.5	430,600
1973	11.5	23.9	64.6	330,500
1974	14.1	63.9	22.0	1,708,800
1975	14.5	34.4	51.1	1,270,100
1976	12.7	33.5	53.8	817,000
1977	11.3	39.5	49.2	561,800
1978	14.2	51.3	34.5	2,267,200
1979	7.3	60.4	32.3	1,706,400
1980	20.8	24.5	54.7	2,969,000
1981	23.0	20.7	56.3	1,233,000
1982	14.0	17.2	68.8	976,400
1983	14.3	60.9	24.8	1,361,000
1984	11.4	27.6	61.0	1,002,800
1985	18.6	22.2	59.1	939,000
1986	16.1	23.3	60.6	819,000
1987	27.6	56.1	16.3	1,337,000
1988	31.0	44.4	24.6	866,800
1989	19.6	28.9	51.5	1,186,400
1990				1,069,400
1991			19.0	1,159,900
1992	24.9	56.7	18.4	1,286,300
1993	40.9	34.1	25.0	1,176,100
1994	25.5	36.4	38.1	1,471,900
1995	33.5	52.9	13.6	1,482,200
1996	25.8	39.3	34.9	1,649,600
Mean	19.1	43.5	36.9	1,134,266
1997	9.1	35.3	55.6	1,512,400

¹ Estimated from Wood River tower counts. Rounded to the nearest hundred.

Appendix Table 23. Total escapement estimates of pink salmon, Nushagak and Togiak Districts, 1962-1997.^a

Year	Nushagak District ¹	Togiak District ²
1962	543,000	
1964	910,560	
1974	585,520	8,620 ^d
1976	863,430	37,570
1978	9,386,480	150,000 ^d
1980	2,785,200	102,820
1982	1,656,660	44,300
1984	2,926,450	269,950
1986	72,190 ^b	80,000 ^d
1988	494,610 ^b	142,500 ^d
1990	801,730 ^b	207,000
1992	^c	235,000 ^d
1994	192,780 ^b	88,000 ^d
1996	821,312 ^b	^c
Mean	1,768,218	124,160
1997	non-pink year, even years only	

¹ Includes Wood, Igushik, Snake, Nushagak, and Nuyakuk Rivers, and Ice, Youth, and Sunshine Creeks, unless otherwise noted.

² Includes Togiak, Matogak and Osviak Rivers; 1982 and 1990 also Include Slug River.

^a Only those years of comprehensive aerial coverage are included: even years only; all counts rounded to the nearest 10 fish.

^b Sonar estimate of Nushagak-Mulchatna Rivers only.

^c No escapement estimate.

^d Togiak River estimate only.

Appendix Table 24. Aerial estimates of sockeye salmon escapements, Togiak District, 1977 - 1997.^a

Year	Togiak River & Tributaries ¹	Kulukak Systems ²
1974	20,600	4,900
1975	19,600	8,600
1977	15,600	40,100
1978	30,600	33,900
1979	23,700	26,600
1980	50,700	45,700
1981	39,700	58,800
1982	25,300	52,800
1983	13,200	27,000
1984	30,900	49,800
1985	8,800	36,600
1986	35,000	42,800
1987	28,600	37,800
1988	32,400	31,700
1989	19,800	10,800
1990	47,100	49,600
1991	23,700	23,900
1992	16,500	26,400
1993	15,900	31,800
1994	19,420	29,740
1995	25,508	14,620
1996	30,220	18,980
<hr/>		
1977-96 Mean (20-Year)	26,632	34,472
1977-86 Mean (10-Year)	27,350	41,410
1987-96 Mean (10-Year)	25,915	27,534
<hr/>		
1997	20,625	7,950

¹ Estimates do not include fish spawning above the counting tower (Togiak Lake outlet); estimates for Ungalikthluk, Osviak,. Matogak and Slug Rivers are not included in the 1977-94 data as reported in Bristol Bay Data Reports 73 and 81.

² Includes Kulukak River, Kulukak Lake, and Tithe Creek Ponds.

^a All counts are rounded to the nearest hundred.

Appendix Table 25. Peak aerial counts of live sockeye salmon, Togiak River drainage, 1977 - 1997.

Year	Togiak Mainstem	Gechiak River	Pungokebuk River	Nayorurun River	Kemuk River	Ongivunuck River	Total
1977	2,200	500	2,000			3,100	7,800
1978	10,000	2,020	1,200			4,620	17,840
1979	7,100	520	750			2,800	11,170
1980	18,600	3,200	2,500	500	3,200	2,000	30,000
1981	14,100	2,700	3,150			3,400	23,350
1982	2,300	3,600	2,500	0	100	4,800	13,300
1983	4,800	1,100	700	0	0	1,200	7,800
1984	10,550	2,800	2,450	0	0	2,300	18,100
1985	1,800	400	500	0	0	1,700	4,400
1986	13,500						13,500
1987	5,200	3,600	600	0	0	4,900	14,300
1988	9,400	2,000	1,100	0	0	3,700	16,200
1989	7,600	1,500	630			150	9,880
1990	8,770	5,720	5,980	0	2,550	1,190	24,210
1991	7,990	1,640	1,220			1,010	11,860
1992	3,030	1,280	1,400			2,200	7,910
1993	2,300	1,270	540			2,950	7,060
1994	3,100	560	1,870			3,900	9,430
1995	3,260	1,745	1,000		4,200	2,330	12,535
1996	9,160	2,270	150	100	240	3,190	15,110
Mean	7,238	2,022	1,592	67	1,029	2,707	14,655 ^a
%	49.4%	13.8%	10.9%	0.5%	7.0%	18.5%	100.0%
1997	8,200	1,600	450	50	650	2,800	13,750

^a Sum of means for all streams.

Appendix Table 26. Peak aerial counts of live sockeye salmon, Togiak District, 1977-1997.

Year	Togiak River ¹	Kulukak River ²	Tithe Creek Ponds	Quigmy River	Matogak River	Osviak River	Slug River	Negukthlik River	Ungalikthluk River	Total
1977	7,800	6,400	18,200		200	2,000	2,700		1,700	39,000
1978	17,840	8,100	11,800						1,000	38,740
1979	11,170	4,600	10,800		200	200		600	700	28,270
1980	30,000	12,200	14,200		500	200	1,900			63,500
1981	23,350	15,700	18,250		700	6,400	5,900	3,900	12,800	87,000 ^a
1982	13,300	11,900	19,300		0	1,000	5,500	300	2,400	53,700
1983	7,800	8,430	2,720		80	20	2,000	230	940	22,220
1984	18,100	7,400	14,000		200	6,800		100	5,200	51,800
1985	4,400	6,700	11,600		0	200	2,300	260	1,310	26,770
1986	13,500	10,900	14,000							38,400
1987	14,300	10,500	8,400							33,200
1988	16,200	12,600	3,250	250	100	380	5,880	200	2,700	41,560
1989	9,880	2,920	2,500					5,000		20,300
1990	24,210	10,600	14,200	100	400	2,200	3,540	9,700	3,800	68,750
1991	11,860	8,650	3,320	35	860	2,530	560	3,400	2,650	33,865
1992	7,910	7,530	4,950	40	300	3,340	1,460	3,600	3,760	32,890
1993	7,060	9,600	6,300					3,100	5,680	31,740
1994	9,430	10,270	4,600	580	990	1,750	6,070	2,230	3,240	39,160
1995	12,535	3,000	4,310	200	610	1,470	2,820	390	1,720	27,055
1996	15,110	2,490	7,000 ^b		360	780	1,045	1,000 ^b		27,785
Mean	13,788	8,525	9,685	172	367	1,951	3,206	2,267	3,100	43,060
%	32.0%	19.8%	22.5%	0.4%	0.9%	4.5%	7.4%	5.3%	7.2%	100.0% ^c
1997	13,750	2,300	3,000		360	780	1,045	1,000		22,235 ^b

¹ Includes all surveyed sections of Togiak River proper and all tributaries to the Togiak River.

² Includes surveys of Kulukak Lake. Counts prior to 1977 include Kulukak Lake only and are not included in the mean.

^a Includes a combined count for the Negukthlik and Ungalikthluk of 4,500 fish.

^b Complete count not available.

^c Sum of means for all streams.

Appendix Table 27. Peak aerial counts of live chinook salmon, Togiak River drainage, 1977-1997.

Year	Togiak River Section ¹						Pungokepuk River		Nayorurun River	Kemuk River	Ongivinuck River	Total
	A	B	C	D	E	F	Gechiak River					
1977							1,190	500	230	120	120	2,160
1978	940	1,240	1,390	810	1,060	1,850	2,150	590	780	220	220	11,250
1979	370	250	330	150	560	890	1,060	360	250	170	220	4,610
1980	180	120	340	230	120	140	910	200	510	170	190	3,110
1981	420	390	500	200	300	740	980	310	370	390	290	4,890
1982					80	320	470	170	190	130	470	1,830
1983	120	220	370	290	360	850	820	240	340	430	350	4,390
1984	250	560	900	560	820	1,920	760	580	270	580	430	7,630
1985	270	320	640	340	470	970	470	250	290	310	460	4,790
1986	150	80	160	30	110	350						880
1987	20	70	170	120	200	480	610	180	100	120	320	2,390
1988	70	70	160	160	170	710	390	180	60	70	90	2,130
1989	10	30	370			940	190	80			40	1,660
1990	230	170	680	365	805	1,085	370	125	75	400	10	4,315
1991	505	165	475	225	520	455	460	105	90	100	150	3,250
1992	150	250	440	225	450	690	250	160	70	175	105	2,965
1993	170	120	220	160		1,810 ^a	595	240	130	65	440	3,950
1994				215	815	1,580	420	215	225	570	380	4,420
1995	120	220	750	255	800	800	715	140	425	520	295	5,040
1996	75	150	160	100	255	625	335	120	120	235	325	2,500
Mean	238	260	474	261	464	906	692	250	251	265	258	4,320
%	5.5%	6.0%	11.0%	6.0%	10.8%	21.0%	16.0%	5.8%	5.8%	6.1%	6.0%	100.0% ^b
1997	100	350	1300	600	820	1000	275	180	150	275	100	5150

¹ Section A; Togiak Bay - Gechiak River
Section B; Gechiak River - Pungokepuk River
Section C; Pungokepuk River - Nayorurun River
Section D; Nayorurun River - Kemuk River
Section E; Kemuk River - Ongivinuck River
Section F; Ongivinuck River - Togiak Lake

^a Includes count for Section E.

^b Sum of means for all streams.

Appendix Table 28. Peak aerial counts of live chinook salmon, Togiak District, 1977-1997.

Year	Togiak River ¹	Quigny River	Kulukak River	Matogak River	Osviak River	Slug River	Negukthlik River	Ungalikthluk River	Total
1977	2,160		1,480	60	120		440	40	4,300
1978	11,250		2,720	150	250		1,020	110	15,500
1979	4,610	20	2,260	100	210		850	130	8,180
1980	3,110	0	700	70	40		260	160	4,340
1981	4,890	0	1,290	470	1,730	350	1,460	180	10,370
1982	1,830	90	1,690	290	320		1,600	280	6,100
1983	4,390	40	2,460	190	120		1,080	260	8,540
1984	7,630	30	1,190	150	360		680	20	10,060
1985	4,790	0	540	100	50		80	90	5,650
1986	880								880
1987	2,390		300	30	40		660	80	3,500
1988	2,130	10	490	0	40	0	650	170	3,490
1989	1,660		740				560		2,960
1990	4,315	30	635	75	60	0	930	25	6,070
1991	3,250	25	285	75	100		1,175	55	4,965
1992	2,965	15	485	40	105	30	490	35	4,165
1993	3,950		1,140	80	110	100	830	70	6,280
1994	4,420	20	835	40	60	10	540	190	6,115
1995	5,040	35	430	65	135	50	740	80	6,575
1996	2,500	35	698	35	71	30	402 ^a		3,771 ^a
Mean	3,908	25	1,072	112	218	71	760	110	6,276
%	62.3 %	0.4 %	17.1 %	1.8 %	3.5 %	1.1 %	12.1 %	1.7 %	100.0 % ^b
1997	5,150	10	310	50	65	33		10	5,628

¹ Includes all surveyed sections of Togiak River proper and all tributaries to the Togiak River.

^a Complete count not available.

^b Sum of means for all streams.

Appendix Table 29. Peak aerial counts of live chum salmon, Togiak River drainage, 1977-1997.

Year	Togiak River Section ¹						Gechiak River	Pungokepuk River	Nayorurun River	Kemuk River	Ongivinuck River	Total
	A	B	C	D	E	F						
1977	12,000	8,000	10,900	8,000		15,100	13,600	4,900	22,100	3,100	2,400	100,100
1978	24,500	7,400	7,500	1,600	15,200	3,300	6,300	2,500	7,300	1,800	8,100	85,500
1979	14,000	2,800	3,300	800	6,600	10,400	3,500	1,000	2,500	500	200	45,600
1980	41,300	11,000	9,200	900	6,000	3,100	10,200	2,700	10,100	800	3,500	98,800
1981	11,800	4,500	2,400	1,000	3,000	6,000	3,100	500	4,300	1,700	4,200	42,500
1982				200	1,200	2,500	500	400	1,300	100	1,000	7,200
1983	8,160	3,050	3,780	1,100	2,780	6,070	150	140	5,560	570	3,790	35,150
1984	3,900	6,300	800	0	2,600	6,400	3,700	2,000	4,200	700	3,500	34,100
1985	8,300	6,500	3,200	900	6,700	10,200	4,100	600	9,600	1,800	8,300	60,200
1986 ^a												
1987	12,000	9,400	2,700	500	13,200	33,000	2,600	1,200	4,100	700	13,100	92,500
1988	10,000				4,900	3,800	3,700	5,000	3,500	200	3,800	34,900
1989		2,600	2,100		5,000	8,100	290	700			1,200	19,990
1990	2,200	1,275	1,350	400	650	4,200	3,150	1,150	3,400	250	125	18,150
1991	10,200	3,900	2,800	600	5,500	6,000	2,300	500	3,500	800	3,480	39,580
1992 ^b	1,800	1,800	300	100	1,200	1,500	2,000	500	1,800	900	800	22,700
1993	6,500	3,500	2,300	60		4,400 ^d	1,950	450	4,380	620	3,500	23,260 ^c
1994				1,300	5,200	10,400	900	2,400	7,100	900	5,700	33,900
1995	15,700	7,100	4,700	1,800	6,800	5,900	4,800	1,900	9,700	2,700	8,200	69,300
1996	3,700	10,250	5,500	1,300	5,750	8,250	2,600	750	900	550	3,400	42,950
Mean	12,284	5,838	4,169	1,279	5,693	7,929	3,747	1,599	6,091	1,046	4,305	53,980
%	284.4%	135.2%	96.5%	29.6%	131.8%	183.6%	86.7%	37.0%	141.0%	24.2%	99.7%	100.0% ^e
1997	3,900	3,100	3,800	2,750	7,100	4,550	3,200	800	4,750	1,800	3,900	39,650

¹ Section A; Togiak Bay - Gechiak River
Section B; Gechiak River - Pungokepuk River
Section C; Pungokepuk River - Nayorurun River
Section D; Nayorurun River - Kemuk River
Section E; Kemuk River - Ongivinuck River
Section F; Ongivinuck River - Togiak Lake

^a No aerial surveys conducted.

^b Counts by section are not representative due to post-peak survey, and are not included in the mean.

^c Preferred total estimate; management survey count conducted 7/15/92.

^d Includes count for Section E.

^e Sum of means for all streams.

Appendix Table 30. Peak aerial counts of live chum salmon, Togiak District, 1977-1997.

Year	Togiak River ¹	Quigmy River	Kulukak River	Matogak River	Osviak River	Slug River	Negukthlik River	Ungalikthluk River	Total
1977	100,100	5,800	21,300	15,300	31,200	2,800	20,000	20,500	217,000
1978	85,500	9,400	24,200	15,000	17,500	6,400	7,600	8,000	173,600
1979	45,600	11,000	16,400	13,400	36,200	4,000	3,800	6,600	137,000
1980	98,800	2,700	27,300	5,700	29,500	6,700	18,500	15,000	204,200
1981	42,500	10,800	11,200	21,700	53,000	3,900	3,800	14,600	161,500
1982	7,200	1,300	8,300	3,100	5,500	2,400	160	1,270	29,230
1983	35,150	4,900	12,960	7,600	11,900	1,210	300	7,360	81,380
1984	34,100	6,300	8,500	10,200	18,400		2,100	3,000	82,600
1985	60,200	1,800	7,800	2,860	5,460	8,800	130	14,650	101,700
1986 ^a									
1987	92,500	1,500	22,000	2,300	2,160				120,460
1988	34,900	10,800	35,000	12,000	17,400	7,600	400	11,300	129,400
1989	19,990	2,820	5,580	7,450	4,900		560		41,300
1990	18,150	555	5,550	1,475	2,300	3,650	750	1,300	33,730
1991	39,580	4,420	9,540	4,730	8,700		120	3,020	70,110
1992	22,700 ^b	600	4,800 ^b	4,400	7,100	1,700	100	4,000	45,400
1993	27,660		6,950	1,970	1,360	3,060	20	4,020	45,040
1994	33,900	890	10,700	1,630	2,000	4,360	230	1,090	54,800
1995	138,600	2,200	7,600	5,200	13,920	6,440	1,000	7,200	182,160
1996	42,950	960	7,560	560	810	2,670	40	^c	55,550
Mean	51,583	4,375	13,328	7,188	14,174	4,379	3,312	7,230	105,570
%	48.9%	4.1%	12.6%	6.8%	13.4%	4.1%	3.1%	6.8%	100.0% ^d
1997	39,650	1,700	4,550	3,000	2,500	1,890	^a	^a	53,290

¹ Includes all surveyed sections of Togiak River proper and tributaries to the Togiak River.

^a No aerial surveys conducted.

^b Preferred estimate from a management survey due to post-peak spawning ground survey.

^c Complete count not available.

^d Sum of means for all streams.

Appendix Table 31. Peak aerial counts of live coho salmon, Togiak River drainage, 1980-1997.

Year	Togiak River Section ¹						Gechiak River	Pungokepuk River	Nayorurun River	Kemuk River	Ongivinuck River	Total
	A	B	C	D	E	F						
1980	3,620	1,010	1,740	1,270	5,080	1,860	3,460	760	1,310	860	740	21,710
1981	9,280	580	100	800	370	750	520	360	230	210	1,300	14,500
1982	2,200	1,500	150	100	1,400	1,700	1,930	1,740	510	200	11,870	23,300
1983 ^a												
1984	1,440	1,190	200	120	620	1,480	4,750	2,240	990	1,110	6,140	20,280
1985	800 ^b	660 ^b	110 ^b	70 ^b	150	820	1,340	750	40	80	6,250	9,430
1986			60	400	100	400					2,560	3,520
1987	340	500	250	200	240	530	1,020	70			1,060	4,210
1988	950	370		140	210	360	1,530				4,100	8,590
1989 ^a												
1990	1,650	390	400	0	540	660	920	450	260	130	1,730	7,130
1991	4,900 ^c	400 ^c	700 ^c	600 ^c	1,680 ^c	140					100 ^c	140 ^c
1992	4,420	1,120	1,180	540	2,940	3,080	5,240	1,440	780	1,500	4,460	26,700
1993 ^a												
1994 ^a							1,290 ^c	220 ^c	120 ^c	95 ^c	1,930	
1995 ^a							1,450			200	1,180	
1996	2,550	1,090	150	250	1,600	5,020	2,080	1,170	575	725	6,450	21,660
Mean	2,645	775	423	347	1,205	1,515	2,020	898	522	502	3,828	14,679 ^d
%	61.2%	17.9%	9.8%	8.0%	27.9%	35.1%	46.8%	20.8%	12.1%	11.6%	88.6%	100.0%
1997	600	200	400	100	400	1,800	1,000	650	350	475	900	6,875

¹ Section A; Togiak Bay - Gechiak River
Section B; Gechiak River - Pungokepuk River
Section C; Pungokepuk River - Nayorurun River
Section D; Nayorurun River - Kemuk River
Section E; Kemuk River - Ongivinuck River
Section F; Ongivinuck River - Togiak Lake

^a No aerial surveys conducted.

^b Proportional estimates based on 1984 data.

^c Timing of aerial surveys did not coincide with the period of peak spawning activity, and therefore, counts were not included in the mean or percent.

^d Sum of means for all streams.

Appendix Table 32. Peak aerial counts of live coho salmon, Togiak District, 1980-1997.

Year	Togiak River ¹	Quigmy River	Kulukak River	Matogak River	Osviak River	Slug River	Negukthlik River	Ungalikthluk River	Creek	Total
1980	21,710		10,300							32,010
1981	14,500		3,790				100	840	1,080	20,310
1982	23,300		3,380							26,680
1983 ^a										
1984	20,280		10,750	1,850	1,080	670				34,630
1985	9,430	200	7,790	610	420					18,450
1986	3,520									3,520
1987	4,210	30	910	440	120			130		5,840
1988	8,590	460	1,840	310	490	470	370	3,170		15,700
1989 ^a										
1990	7,130	1,029	5,195	2,675	1,491	810		4,153		22,483
1991	140 ^b		4,200 ^b							0
1992	26,700		12,640							39,340
1993 ^a										
1994 ^a										
1995		855	1,185	1,392	1,080	1,149		5,196 ^c		
1996	21660	1,211	10,290	3,062	2,805	1,944	851	5917	^f	47740
Mean	13,419	631	5,673	1,477	1,069	1,009	440	2,368	540	26,626 ^d
%	50.4%	2.4%	21.3%	5.5%	4.0%	3.8%	1.7%	8.9%	2.0%	100.0%
1997	6,875	325	1,675	150	1,046	1,397		1,690		13,158

¹ Includes all surveyed sections of Togiak River proper and tributaries to the Togiak River. See Appendix Table 34.

^a No aerial surveys conducted.

^b Timing of aerial surveys did not coincide with the period of peak spawning activity, and therefore, counts were not included in the mean or percent.

^c Only Togiak River tributaries surveyed; not included in the mean or percent.

^d Sum of means for all streams.

^e Negukthlik and Ungalikthluk Rivers combined.

^f Complete count not available.

The Alaska Department of Fish and Game administers all programs and activities free from discrimination on the basis of sex, color, race, national origin, age, marital status, pregnancy, parenthood, or disability. For information on alternative formats available for this and other department publications, contact the department ADA Coordinator at (voice) 907-465-4120, or (TDD) 907-465-3646. Any person who believes s/he has been discriminated against should write to: ADF&G, P.O. Box 25526, Juneau, AK 99802-5526; or O.E.O.

